

Characterization of the Consciousness Energy Healing Treated Essential Amino Acid: L-Tryptophan

Branton A¹, Trivedi MK¹, Trivedi D¹, Nayak G¹ and Jana S^{*2}

¹Trivedi Global, Inc., Henderson, USA

²Trivedi Science Research Laboratory Pvt. Ltd., Thane (W), India

*Corresponding author: Jana S, Trivedi Science Research Laboratory Pvt. Ltd., Thane (W), Maharashtra, India, Tel: +91- 022-25811234, Email: publication@trivedieffect.com

Citation: Branton A, Trivedi MK, Trivedi D, Nayak G, Jana S (2019) Characterization of the Consciousness Energy Healing Treated Essential Amino Acid: L-Tryptophan. J Adv Food Technol 2(1): 102

Received Date: July 22, 2019 **Accepted Date:** October 01, 2019 **Published Date:** October 03, 2019

Abstract

L-tryptophan is an amino acid which helps in the formation of proteins and certain brain-signaling chemicals, etc. This study was carried out to investigate the impact of the Trivedi Effect® on the physicochemical properties of L-tryptophan by using modern analytical techniques. For analysis, the sample was divided into control and treated parts. In which control part didn't received any Biofield Energy Treatment, while the treated part was received the Trivedi Effect®-Consciousness Energy Healing Treatment was remotely by a famous Biofield Energy Healer, Alice Branton. The values of particle size in the treated sample was increased by 1.52% (d_{10}), 1.24% (d_{50}), 4.33% (d_{90}), and 3.31% {D(4,3)}, while the specific surface area was reduced by 1.05% as compared to the control sample. The treated L-tryptophan shown that the powder XRD peak intensities and crystallite sizes were altered, ranging from -87.32% to 20.88% and -9.54% to 614.29%, respectively, compared to the control sample. Moreover, the average crystallite size of the treated L-tryptophan was increased by 25.97% than the control sample. The overall weight loss of the treated L-tryptophan was reduced by 5.27%, in comparison to the control sample. It resulted in a significant 228.88% increase in the residual amount of the treated L-tryptophan, as compared to the untreated sample. Besides, the DTG thermograms of both the samples showed two peaks and the maximum thermal degradation temperature corresponding to 1st and 2nd peak of the treated sample was significantly altered by 25.34% and -1.45%, respectively as compared to the control sample. The melting temperature of the treated sample was increased by 2.69%, while the ΔH_{fusion} was reduced by 16.36%, as compared with the control sample. The Consciousness Energy Healing Treatment might act in creating some novel polymorphic form of L-tryptophan that might prove to be beneficial in terms of improved flowability, appearance, and thermal stability as compared to the untreated sample. Thus, the use of the Biofield Energy Treated L-tryptophan might be considered as an advantageous approach for designing the more efficacious nutraceutical/pharmaceutical formulations.

Key words: The Trivedi Effect®; Consciousness Energy Healing Treatment; L-tryptophan; Particle size; PXRD; TGA/DTG

Introduction

L-tryptophan is one of the essential amino acids used in the biosynthesis of proteins and certain brain-signaling chemicals. It is as a precursor for serotonin, which is a neurotransmitter that helps in controlling the mood and sleep, and synthesized it with the help of tryptophan hydroxylase [1]. L-tryptophan is metabolized further into melatonin and vitamin B₃ (nicotinic acid), in which the melatonin plays a vital role in maintaining the circadian rhythms of various biological functions [2]. L-tryptophan can be obtained naturally from animal and plant proteins. It plays a significant role in the development as well as the functioning of various organs of the body. The body absorbs L-tryptophan from food and converts it in the form of 5-HTP (5-hydroxytryptophan), followed by serotonin, melatonin, and vitamin B₆ (nicotinamide) [3,4]. Serotonin helps in transmitting the signals between the nerve cells and also causes the narrowing of blood vessels [5]. The reduced levels of tryptophan inside the body may cause depression, obsessive-compulsive disorder, and bipolar disorder, etc. [6]. It is used for the treatment of some mental health disorders [7], for improving athletic performance [8]; quit smoking [9], treatment of emotional symptoms in premenstrual dysphoric disorder (PMDD) [10], attention deficit-hyperactivity disorder (ADHD), mental functionality problems in the elderly [11], ulcer healing caused by H pylori [12], sleep apnea, seasonal affective disorder (SAD), and anxiety [13], etc. It is soluble in water (11.4g/L at 25 °C), hot alcohol, and alkali hydroxide; insoluble in chloroform.

The physicochemical properties of any nutraceutical have very much importance in its quality. Biofield Energy Treatment has grate role altering the physicochemical properties of nutraceutical compounds [14]. Biofield is a putative energy field, which has the basic concept that human beings possess this subtle form of energy [15]. Thus, an expert has the ability to harness energy from the universe and can communicate it to any object(s) around the globe. Such energy is considered as the Biofield Energy, and its healing techniques are

known as the Biofield Energy Treatment, which was also accepted by the National Center for Complementary and Alternative Medicine (NCCAM) [16] under CAM. The other therapies involved under this category are deep breathing, yoga, Tai Chi, Qi Gong, meditation, chiropractic/osteopathic manipulation, massage, special diets, homeopathy, progressive relaxation, acupressure, guided imagery, acupuncture, relaxation techniques, hypnotherapy, movement therapy, healing touch, pilates, rolfing structural integration, traditional Chinese herbs and medicines, mindfulness, Ayurvedic medicine, naturopathy, essential oils, Reiki, aromatherapy, cranial-sacral therapy and applied prayer [17,18]. The Trivedi Effect®-Consciousness Energy Healing Treatment has also been considered worldwide due to its remarkable results in the living organisms as well as the non-living materials. Biofield Energy Healing Treatment has been reported for causing significant alterations in the physicochemical and thermal properties of various organic/pharmaceutical compounds [19-22], alter the characteristics of crops [23,24], microorganisms [25,26], metals, ceramics, and polymers [27-29], nutraceuticals [14], and improved the bone and skin health [30-32], etc. Thus, the aim of this study was to analyse the effect of the Biofield Energy Healing Treatment on the physicochemical and thermal properties of L-tryptophan by using various analytical techniques.

Materials and Methods

Chemicals and Reagents

The test sample L-tryptophan was purchased from Alfa Aesar, India. Similarly, other chemicals used in the experiments also purchased in India.

Consciousness Energy Healing Treatment Strategies

The L-tryptophan sample considered for the experiment was divided into two equal parts. One part of L-tryptophan sample was considered as a control sample where no Biofield Energy Treatment was provided. However, the other part of L-tryptophan was received the Trivedi Effect®-Consciousness Energy Healing Treatment distantly under typical laboratory conditions (3 minutes) by the famous Biofield Energy Healer, Alice Branton, USA, known as the treated L-tryptophan sample. However, the control sample was treated under same laboratory conditions with a “sham” healer who is ignorant about the Biofield Energy Treatment. After that, both the samples were kept in sealed conditions and characterized using sophisticated analytical techniques.

Characterization

The particle size distribution (PSD) analysis of L-tryptophan was performed using Malvern Mastersizer 2000 (UK) using the wet method [33,34]. The X-ray diffraction (PXRD) analysis of L-tryptophan powder sample was performed with the help of Rigaku MiniFlex-II Desktop X-ray diffractometer (Japan) [35,36]. The crystallites size was calculated with the help of the Scherrer's formula (1)

$$G = k\lambda / \beta \cos \theta \quad (1)$$

Where G = crystallite size (nm), λ = radiation wavelength, k = equipment constant, β = full-width at half maximum, and θ = Bragg angle [37].

The thermal gravimetric analysis (TGA) thermograms of L-tryptophan were obtained with the help of TGA Q50 TA instruments. The differential scanning calorimetry (DSC) analysis of L-tryptophan was performed with the help of DSC Q200, TA instruments [38].

The % change in the above parameters the treated L-tryptophan was calculated compared to the control sample using the following equation 2:

$$\% \text{ change} = \frac{[\text{Treated} - \text{Control}]}{\text{Control}} \times 100 \quad (2)$$

Results and Discussion

Particle Size Analysis (PSA)

The analysis revealed the particle size distribution of the control and Biofield Energy Treated L-tryptophan samples in Table 1. It showed the particle size distribution of the control sample at $d_{10} = 27.61 \mu\text{m}$, $d_{50} = 140.51 \mu\text{m}$, $d_{90} = 364.34 \mu\text{m}$, and $D(4, 3) = 172.08 \mu\text{m}$. Besides, the particle size distribution of the Biofield Energy Treated sample showed significant alterations as it was observed at $d_{10} = 28.03 \mu\text{m}$, $d_{50} = 142.25 \mu\text{m}$, $d_{90} = 380.13 \mu\text{m}$, and $D(4, 3) = 177.77 \mu\text{m}$. Therefore, the analysis showed a significant increase in the particle size values 1.52% (d_{10}), 1.24% (d_{50}), 4.33% (d_{90}), and 3.31% [$D(4,3)$] of the Biofield Energy Treated L-tryptophan compared to the control sample.

Parameter	d_{10} (μm)	d_{50} (μm)	d_{90} (μm)	$D(4,3)$ (μm)	SSA(m^2/g)
Control	27.61	140.51	364.34	172.08	0.095
Biofield Energy Treated	28.03	142.25	380.13	177.77	0.094
Percent change (%)	1.52	1.24	4.33	3.31	-1.05

d_{10} , d_{50} , and d_{90} : particle diameter corresponding to 10%, 50%, and 90% of the cumulative distribution, $D(4,3)$: average mass-volume diameter, and SSA: specific surface area.

Table 1: Particle size data of the control and treated L-tryptophan

On a similar note, the Biofield Energy Treated sample showed a decrease (1.05%) in the specific surface area as it was observed as 0.094 m²/g, as compared with the SSA of the control sample that was found as 0.095 m²/g. This significant alteration in the surface area of the Biofield Energy Treated L-tryptophan may arise because of the increase in particle size after the Biofield Energy Treatment. Later on, the particle size distribution of the drug is known as an important factor in the formulation design as it might impact the flowability, blend uniformity, and compactibility factors of the drug with excipients. Such properties are also considered important in defining the safety, efficacy, and the quality of the drug formulation. The literature reported that the flowability of the drug might be hindered in the presence of smaller particles as compared to the large and spherical particles [39,40]. Hence, it is presumed that the treated sample might show better flowability, uniformity, and compactibility during the formulation development as compared to the untreated sample.

Powder X-ray Diffraction (PXRD) Analysis

The PXRD diffractograms of the control and treated L-tryptophan samples are shown in Figure 1. Moreover, the calculations corresponding to the peak intensities and crystallite sizes and further analysis for both the samples are presented in Table 2.




Figure 1: PXRD diffractograms of the control and treated L-tryptophan

Entry No.	Bragg angle (°2θ)		Intensity (cps)			Crystallite size (G, nm)		
	Control	Treated	Control	Treated	% change	Control	Treated	% change
1	4.79	4.81	2491	3011	20.88	227.0	267.0	17.62
2	9.61	9.64	769	549	-28.61	208.0	239.0	14.90
3	14.53	14.54	2273	1574	-30.75	222.0	226.0	1.80
4	18.24	18.32	375	57	-84.80	81.0	104.0	28.40
5	19.51	19.71	593	299	-49.58	233.0	299.0	28.33
6	24.52	24.74	615	78	-87.32	30.8	220.0	614.29
7	34.78	34.87	116	92	-20.69	262.0	237.0	-9.54

Table 2: PXRD data for the control and treated L-tryptophan

The diffractogram analysis revealed the alteration in the Bragg's angles of the characteristic peaks of the Biofield Energy Treated sample as compared to the control L-tryptophan sample. Besides, the peak intensities of the Biofield Energy Treated sample also showed significant changes ranging from -87.32% to 20.88% in comparison to the control sample. Similarly, the crystallite sizes of the Biofield Energy Treated sample were also observed to be changed, ranging from -9.54% to 614.29%, as compared to the control sample. Besides, the average crystallite size of the control sample was observed to as 180.54 nm; while the Biofield Energy Treated sample showed an average crystallite size as 227.43 nm. Thus, the average crystallite size of the treated sample was significantly increased by 25.97% as compared to the control sample. The Biofield Energy Treatment was reported by several kind of literatures regarding its possible impact on the compound to alter the crystalline structure and crystal morphology by altering the peak intensities and crystallite sizes, and thereby might create a new polymorph of the compound [41]. Thus, the analysis indicated the formation of the new polymorph of the compound through significant alterations in the peak intensities, and crystallite size of the treated L-tryptophan, which could be used in improving its drug profile [42] as compared to the untreated sample.

Thermal Gravimetric Analysis (TGA)/ Differential Thermogravimetric Analysis (DTG)

The thermal analysis of both the L-tryptophan samples was done using the TGA/DTG technique. The literature reported the thermogravimetric records for L-tryptophan, according to which it shows a fast mass loss of 20-24% starting at 526-538 K, followed by a continuous mass decrease. Also, it was reported that some small molecules of NH_3 , CO_2 , and H_2O were evolved during the degradation [43]. The TGA thermograms of both the samples (Figure 2) showed thermal degradation at $\sim 250^\circ\text{C}$, which was similar as reported in the literature. The further analysis revealed that the total weight loss of the control sample was 97.75%, while it was observed as 92.60% for the Biofield Energy Treated sample. Thus, the total weight loss of the treated L-tryptophan sample was reduced by 5.27% during the sample degradation as compared to the untreated sample. As a result, the residue amount of the treated sample was significantly increased by 228.88% (Table 3) as compared to the control sample. It indicated the reduction in thermal degradation and improved thermal stability of the Biofield Energy Treated L-tryptophan sample as compared to the untreated sample.




Figure 2: TGA thermograms of the control and treated L-tryptophan

Sample	TGA		DTG	
	Total weight loss (%)	Residue %	Peak 1 T_{\max} ($^\circ\text{C}$)	Peak 2 T_{\max} ($^\circ\text{C}$)
Control	97.75	2.25	228.60	394.34

Sample	TGA		DTG	
	Total weight loss (%)	Residue %	Peak 1 T_{max} (°C)	Peak 2 T_{max} (°C)
Biofield Energy Treated	92.60	7.40	286.53	388.60
% Change	-5.27	228.88	25.34	-1.45

T_{max} = The temperature at which maximum weight loss takes place in TG or peak temperature in DTG.

Table 3: TGA/DTG data of the control and treated L-tryptophan

Moreover, the DTG thermograms of the control and Biofield Energy Treated samples (Figure 3) showed two peaks. The analysis revealed that the maximum degradation temperature (T_{max}) corresponding to 1st peak of the treated sample was increased by 25.34%; while it was slightly decreased by 1.45% corresponding to 2nd peak as compared to the control sample. It indicated that the maximum thermal degradation temperature of the Biofield Energy Treated sample was significantly improved in comparison to the untreated sample. Overall, the TGA/DTG studies showed the improved thermal stability profile of the Biofield Energy Treated L-tryptophan sample as compared with the control sample.




Figure 3: DTG thermograms of the control and treated L-tryptophan

Differential Scanning Calorimetry (DSC) Analysis

The DSC technique helps in studying the thermal behaviour of the compound such as melting and crystallization temperature etc. [44]. Previous literature reported the presence of a single endothermic peak in the thermogram of L-tryptophan, in the temperature range of 540-577 K [45].

The DSC thermograms of both the samples were observed to be similar as reported in the literature and they also showed a single endothermic peak (Figure 4), which is considered as the melting peak. The peak temperature of the control sample was 283.40 °C, while it was observed at 291.02 °C in the treated sample. Thus, the analysis showed that the peak temperature of the Biofield Energy Treated sample was increased by 2.69%, as compared to the control sample (Table 4). Moreover, the latent heat of fusion (ΔH) of the treated sample was observed as 316.50J/g, which was significantly reduced by 16.36% compared with the ΔH of the control sample (378.40J/g).

The analysis indicated the improved thermal stability of the treated L-tryptophan sample after the Biofield Energy Treatment. Thus, it was assumed that the Biofield Energy Treatment cause some alterations in the molecular chains and crystallization structure [46] of the treated L-tryptophan, which might be responsible for such alterations in the melting temperature and latent heat of fusion as compared to the untreated sample.




Figure 4: DSC thermograms of the control and treated L-tryptophan

Sample	Melting point (°C)	ΔH (J/g)
Control Sample	283.40	378.40
Biofield Energy Treated	291.02	316.50
% Change	2.69	-16.36

ΔH: Latent heat of fusion

Table 4: DSC data for the control and treated samples of L-tryptophan

Conclusion

The overall results of the control and Biofield Energy Treated L-tryptophan indicated that the Trivedi Effect®-Consciousness Energy Healing Treatment imposes the significant impact on the physicochemical properties such as particle size, surface area, peak intensities, crystallite size, thermal degradation, melting point, and other thermal properties of the L-tryptophan sample. The particle size values of the Biofield Energy Treated sample was increased by 1.52% (d_{10}), 1.24% (d_{50}), 4.33% (d_{90}), and 3.31% {D(4,3)} compared with the control sample. Moreover, it also resulted in 1.05% decrease in the specific surface area of the Biofield Energy Treated sample, as compared to the untreated L-tryptophan sample. The peak intensities of the Biofield Energy Treated sample were observed to be altered ranging from -87.32% to 20.88%, while the crystallite sizes were altered ranging from -9.54% to 614.29%, compared to the untreated sample. Also, the average crystallite size the Biofield Energy Treated L-tryptophan sample was found to be increased by 25.97% as compared with the control sample. The total weight loss of the treated sample was reduced by 5.27%, compared to the control sample. It resulted in a significant 228.88% increase in the residual amount of the treated L-tryptophan, as compared to the control sample. It revealed the improved thermal stability of the treated sample when compared with the untreated sample. Later on, the DTG data revealed two peaks in the thermograms of both the samples. The analysis showed that the T_{max} corresponding to 1st peak of the Biofield Energy Treated sample was significantly increased by 25.34%; whereas, the

T_{max} corresponding to 2nd peak was slightly decreased by 1.45% as compared to the control sample. The melting temperature of the treated sample was increased by 2.69%, while the ΔH_{fusion} was reduced by 16.36%, as compared to the control sample. It also indicated the improved thermal stability of the Biofield Energy Treated L-tryptophan as compared with the untreated sample. The overall analysis concluded that the Trivedi Effect®-Consciousness Energy Healing Treatment imposes a major impact on the physicochemical and thermal properties of the treated L-tryptophan sample that might improve the flowability, appearance, thermal stability, and storage properties of the sample, compared with the untreated sample. Hence, the Biofield Energy Treated L-tryptophan may be considered suitable for formulation of pharmaceuticals/nutraceuticals regarding the prevention and treatment of various diseases such as mental health disorders, improve athletic performance; treatment of emotional symptoms in premenstrual dysphoric disorder, quit smoking, attention deficit-hyperactivity disorder, ulcer healing caused by *H pylori*, mental functionality problems in the elderly, sleep apnea, seasonal affective disorder, and anxiety, etc.

Acknowledgement

The authors are grateful to Central Leather Research Institute, SIPRA Lab. Ltd., Trivedi Science, Trivedi Global, Inc., Trivedi Testimonials, and Trivedi Master Wellness for their assistance and support during this work.

References

- Sidransky H (2002) Tryptophan: Biochemical and health implications. CRC Series in Modern Nutrition, Florida.
- Ahmad F, Moat AG (1966) Nicotinic acid biosynthesis in prototrophs and tryptophan auxotrophs of *Saccharomyces cerevisiae*. J Biol Chem 241: 775-80.
- Palego L, Betti L, Rossi A, Giannaccini G (2016) Tryptophan biochemistry: Structural, nutritional, metabolic, and medical aspects in humans. J Amino Acids: 8952520.
- Nehete JY, Bhambar RS, Narkhede MR, Gawali SR (2013) Natural proteins: Sources, isolation, characterization and applications. Pharmacogn Rev 7: 107-16.
- Jenkins TA, Nguyen JCD, Polglaze KE, Bertrand PP (2016) Influence of tryptophan and serotonin on mood and cognition with a possible role of the gut-brain axis. Nutrients 8: 56.
- Maron E, Shlik J, Nutt DJ (2008) Tryptophan research in panic disorder. Int J Tryptophan Res 1: 3-12.
- Bell C, Forshall S, Adrover M, Nash J, Hood S, et al. (2002) Does 5-HT restrain panic? A tryptophan depletion study in panic disorder patients recovered on paroxetine. J Psychopharmacol 16: 5-14.
- Williams M (2005) Dietary supplements and sports performance: Amino acids. J Int Soc Sports Nutr 2: 63-7.
- Bowen DJ, Spring B, Fox E (1991) Tryptophan and high-carbohydrate diets as adjuncts to smoking cessation therapy. J Behav Med 14: 97-110.
- Freeman EW, Sondheimer SJ (2003) Premenstrual dysphoric disorder: Recognition and treatment. Prim Care Companion J Clin Psychiatry 5: 30-9.
- Berney A, Nishikawa M, Benkelfat C, Debonnel G, Gobbi G, et al. (2008) An index of 5-HT synthesis changes during early antidepressant treatment: Alpha-[11C]methyl-L-tryptophan PET study. Neurochem Int 52: 701-8.
- Celinski K, Konturek PC, Konturek SJ, Slomka M, Cichoz-Lach H, et al. (2011) Effects of melatonin and tryptophan on healing of gastric and duodenal ulcers with *Helicobacter pylori* infection in humans. J Physiol Pharmacol 62: 521-6.
- Argyropoulos SV, Hood SD, Adrover M, Bell CJ, Rich AS, et al. (2004) Tryptophan depletion reverses the therapeutic effect of selective serotonin reuptake inhibitors in social anxiety disorder. Biol Psychiatry 56: 503-9.
- Trivedi MK, Branton A, Trivedi D, Nayak G, Plikerd WD, et al. (2017) A Systematic study of the biofield energy healing treatment on physicochemical, thermal, structural, and behavioral properties of magnesium gluconate. Int J Bioorg Chem 2: 135-45.
- Berman JD, Straus SE (2004) Implementing a research agenda for complementary and alternative medicine. Annu Rev Med 55: 239-54.
- Hintz KJ, Yount GL, Kadar I, Schwartz G, Hammerschlag R, et al. (2003) Bioenergy definitions and research guidelines. Altern Ther Health Med 9: A13-30.
- Jain S, Hammerschlag R, Mills P, Cohen L, Krieger R, et al. (2015) Clinical studies of biofield therapies: summary, methodological challenges, and recommendations. Glob Adv Health Med 4: 58-66.
- Barnes PM, Bloom B, Nahin RL (2008) Complementary and alternative medicine use among adults and children: United States, 2007. Natl Health Stat Report 12: 1-23.
- Trivedi MK, Branton A, Trivedi D, Nayak G, Panda P, et al. (2016) Evaluation of the isotopic abundance ratio in biofield energy treated resorcinol using gas chromatography-mass spectrometry technique. Pharm Anal Acta 7: 481.
- Trivedi MK, Branton A, Trivedi D, Nayak G, Bairwa K, et al. (2015) Spectroscopic characterization of disodium hydrogen orthophosphate and sodium nitrate after biofield treatment. J Chromatogr Sep Tech 6: 282.
- Trivedi MK, Patil S, Shettigar H, Bairwa K, Jana S (2015) Effect of biofield treatment on spectral properties of paracetamol and piroxicam. Chem Sci J 6: 98.
- Trivedi MK, Tallapragada RM, Branton A, Trivedi D, Nayak G, et al. (2015) Potential impact of biofield treatment on atomic and physical characteristics of magnesium. Vitam Miner 3: 129.
- Trivedi MK, Branton A, Trivedi D, Nayak G, Gangwar M, et al. (2015) Agronomic characteristics, growth analysis, and yield response of biofield treated mustard, cowpea, horse gram, and groundnuts. International Journal of Genetics and Genomics 3: 74-80.
- Trivedi MK, Branton A, Trivedi D, Nayak G, Mondal SC, et al. (2015) Evaluation of plant growth, yield and yield attributes of biofield energy treated mustard (*Brassica juncea*) and chick pea (*Cicerarietinum*) seeds. Agriculture, Forestry and Fisheries 4: 291-5.
- Trivedi MK, Patil S, Shettigar H, Mondal SC, Jana S (2015) Evaluation of biofield modality on viral load of Hepatitis B and C viruses. J Antivir Antiretrovir 7: 083-8.
- Trivedi MK, Branton A, Trivedi D, Nayak G, Charan S, et al. (2015) Phenotyping and 16S rDNA analysis after biofield treatment on *Citrobacter braakii*: A urinary pathogen. J Clin Med Genom 3: 129.
- Trivedi MK, Patil S, Tallapragada RM (2013) Effect of biofield treatment on the physical and thermal characteristics of vanadium pentoxide powders. J Material Sci Eng S11: 001.

28. Trivedi MK, Branton A, Trivedi D, Nayak G, Sethi KK, et al. (2016) Gas chromatography-mass spectrometry based isotopic abundance ratio analysis of biofield energy treated methyl-2-naphthylether (Nerolin). *Am J Phys Chem* 5: 80-6.
29. Trivedi MK, Tallapragada RM, Branton A, Trivedi D, Nayak G, et al. (2015) Characterization of physical and structural properties of aluminum carbide powder: Impact of biofield treatment. *J Aeronaut Aerospace Eng* 4: 142.
30. Koster DA, Trivedi MK, Branton A, Trivedi D, Nayak G, et al. (2018) Evaluation of biofield energy treated vitamin D₃ on bone health parameters in human bone osteosarcoma cells (MG-63). *Biochem and Mol Biol* 3: 6-14.
31. Singh J, Trivedi MK, Branton A, Trivedi D, Nayak G, et al. (2017) Consciousness energy healing treatment based herbomineral formulation: A safe and effective approach for skin health. *Am J of Pharmacol and Phytotherapy* 2: 1-10.
32. Smith DM, Trivedi MK, Branton A, Trivedi D, Nayak G, et al. (2017) Skin protective activity of consciousness energy healing treatment based herbomineral formulation. *J of Food and Nutr Sci* 5: 86-95.
33. Trivedi MK, Sethi KK, Panda P, Jana S (2017) A comprehensive physicochemical, thermal, and spectroscopic characterization of zinc (II) chloride using X-ray diffraction, particle size distribution, differential scanning calorimetry, thermogravimetric analysis/differential thermogravimetric analysis, ultraviolet-visible, and Fourier transform-infrared spectroscopy. *Int J of Pharm Invest* 7: 33-40.
34. Trivedi MK, Sethi KK, Panda P, Jana S (2017) Physicochemical, thermal and spectroscopic characterization of sodium selenate using XRD, PSD, DSC, TGA/DTG, UV-vis, and FT-IR. *Marmara Pharm J* 21: 311-8.
35. Desktop X-ray Diffractometer "MiniFlex+" (1997) *The Rigaku Journal* 14: 29-36.
36. Zhang T, Paluch K, Scalabrino G, Frankish N, Healy AM, et al. (2015) Molecular structure studies of (1S,2S)-2-benzyl-2,3-dihydro-2-(1Hinden-2-yl)-1H-inden-1-ol. *J Mol Struct* 1083: 286-99.
37. Langford JI, Wilson AJC (1978) Scherrer after sixty years: A survey and some new results in the determination of crystallite size. *J Appl Cryst* 11: 102-13.
38. Trivedi MK, Branton A, Trivedi D, Nayak G, Plikerd WD, et al. (2017) A systematic study of the biofield energy healing treatment on physicochemical, thermal, structural, and behavioral properties of iron sulphate. *Int J Bioorg Chem* 2: 135-45.
39. Hlinak AJ, Kuriyan K, Morris KR, Reklaitis GW, Basu PK (2006) Understanding critical material properties for solid dosage form design. *J Pharm Innov* 1: 12-7.
40. Morin G, Briens L (2013) The effect of lubricants on powder flowability for pharmaceutical application. *AAPS Pharm Sci Tech* 14: 1158-68.
41. Trivedi MK, Branton A, Trivedi D, Nayak G, Lee AC, et al. (2017) Evaluation of the impact of biofield energy healing treatment (the Trivedi Effect*) on the physicochemical, thermal, structural, and behavioural properties of magnesium gluconate. *Int J Nutr and Food Sci* 6: 71-82.
42. Savjani KT, Gajjar AK, Savjani JK (2012) Drug Solubility: Importance and Enhancement Techniques. *ISRN Pharm* 2012: 195727.
43. Mello GS, Cardoso AP, Oliveira EWRS, Siqueira AB (2015) Tryptophan. A proposal of the mechanism of thermal decomposition. *J Therm Anal Calorim* 122: 1395-401.
44. Gill P, Moghadam TT, Ranjbar B (2010) Differential scanning calorimetry techniques: Applications in biology and nanoscience. *J Biomol Tech* 21: 167-93.
45. Gheorghe D, Neacsu A, Contineanu I, Tanasescu S, Perisanu S (2017) A calorimetric study of L-, D- and DL-isomers of tryptophan. *J Therm Anal Calorim* 130: 1145-52.
46. Zhao Z, Xie M, Li Y, Chen A, Li G, et al. (2015) Formation of curcumin nanoparticles *via* solution enhanced dispersion by supercritical CO₂. *Int J Nanomedicine* 10: 3171-81.

Submit your next manuscript to Annex Publishers and benefit from:

- ▶ Easy online submission process
- ▶ Rapid peer review process
- ▶ Online article availability soon after acceptance for Publication
- ▶ Open access: articles available free online
- ▶ More accessibility of the articles to the readers/researchers within the field
- ▶ Better discount on subsequent article submission

Submit your manuscript at

<http://www.annexpublishers.com/paper-submission.php>