

Mucoadhesive Drug Delivery Systems- An Overview

Ms. Chavan Mayuri¹, Dr. Pradum Ige^{2*}

¹Pratibhatai Pawar College of Pharmacy, Shrirampur, Dist Ahmednagar 413709

²Abasaheb Kakade College of B.Pharmacy, Bodhegaon, Tal. Shevgaon Dist. Ahmednagar 414503

***Corresponding Author:** Dr. Pradum Ige, Abasaheb Kakade College of B.Pharmacy, Bodhegaon, Tal. Shevgaon Dist. Ahmednagar 414503. Tel: 8668388789 E-mail: pradyumna_1978@rediffmail.com

Citation: Ms. Chavan Mayuri, Dr. Pradum Ige (2023) Mucoadhesive drug delivery systems- An Overview. J Pharma Drug Develop 10(1): 101

Abstract

The term mucoadhesion can be defined as a phenomenon of interfacial molecular attractive forces in the layer of surface of a biological membrane and the natural or synthetic polymer, which allows the polymer to adhere to the surface of that membrane for an extended as well as prolonged period of time. Since the last four decades, the conception of mucoadhesion has achieved an important and precious interest in the various fields of pharmaceuticals. There are many advantages of mucoadhesive buccal drug delivery system that made this a novel drug delivery system for the local as well as systemic delivery of various drugs. The main advantage of this route for drug delivery is that, the drug by this route bypasses the first pass metabolism of various drugs that are liable to their hepatic first pass metabolism. This review provides the brief knowledge about the oral mucosal drug delivery by discussing briefly the structural features of mucosa, mechanism of mucoadhesion, different theories of mucoadhesion, general considerations in the design of mucoadhesive buccal dosage forms, permeation enhancers and the various evaluation methods along with the literature survey of the buccal mucoadhesive drug delivery system.

Keywords: Mucoadhesion; Polyacrylic Acid; Bioadhesion; Bioavailability; Films; Permeation Enhancers; Polymers.

Introduction

Mucoadhesive drug delivery system is a drug delivery system which utilizes the property of bioadhesion of certain polymers which become adhesive on hydration and hence can be used for targeting a drug to a particular region of the body for an extended period of time. Mucoadhesion may be described because the kingdom where substances adhere to every different for a prolonged time frame with the assist of interfacial forces. When this type of substance is organic in nature, the manner is referred to as bioadhesion. Mucoadhesion is the manner of binding a fabric to the mucosal layer of the frame. Utilizing herbal and artificial polymers, mucoadhesive drug delivery is the way of managed drug release which lets for intimate contact between the polymers and the target tissue. Mucoadhesive drug delivery systems are delivery systems which make use of the assets of bioadhesion of positive polymers which grow to be adhesive on hydration and subsequently are able to be used for targeted delivery of the drug to a particular region of the body for a prolonged time period. The concept of mucoadhesion was introduced with the

control released drug delivery within the early 1980s. eight control released system provides continuous drug released at a predetermine rate. In recent years' considerable interest has been shown in the use of bio adhesive polymers and copolymers in controlled drug delivery. This interest is due to the following potential applications of bio adhesive drug delivery system:

- a) Adhesion to specific site of the body, such as the oral and nasal cavities, resulting in an enhanced drug bioavailability.
- b) The formation of an optimum contact with the adhesion surface, increasing drug absorption.
- c) The prolonging of the residence time of the dosage of within ten gastrointestinal tracts. this would reduce the need for multiple dosing, resulting better patient compliance.

The biological surface can be epithelial tissue or the mucus coat on the surface a tissue. If adhesive attachment is to a mucus coat, the phenomenon is referred to as mucoadhesion. Mucoadhesion should not be confused with bio adhesion; in bio adhesion, the polymer is attached to the biological membrane and if the substrate is mucus membrane the term mucoadhesion is used [11]

Advantages of Mucoadhesive Drug Delivery System:

- 1) Drugs shows bypass first pass metabolism then it increases bioavailability.
- 2) Drug is easily administered of therapy in emergency condition.
- 3) Some drugs are not stable in acidic environment of stomach can be administered by buccal delivery.
- 4) Drug release for prolonged period of time.
- 5) In this system drug absorbed by passive diffusion.
- 6) Adaptability in actual state, shape, size & surface [2].

Limitations of Mucoadhesive Drug Delivery System:

- 1) Drug which are not stable at buccal PH cannot be administered.
- 2) Drugs which have bitter taste or not pleasant taste or a good or bother mucosa can't be administered by this route.
- 3) Drug needed with little portion must be controlled.
- 4) Those medications which are consumed by inactive [2]

Historical Development of Mucoadhesive DDS's

since from the last 40 years the idea of mucoadhesion has supplied the awesome soft where in prolonging the residence time as well as controlled released effect of various bioadhesive dosage forms through different mucosal routs .the formulation based on the mucoadhesive drug delivery system have shown enhanced the bioavailablity of many drug .Mucoadhesive drug delivery formulations were introduced in 1947 when gum tragacanth was mix with dental adhesive powder to apply penicillin to the oral mucosa. Over the year's various others polymers for examples sodium alginate, CMS, guar gum, HEC, and tragacanth have been found to exhibit mucoadhesive properties. during the 1980s polyacrylic acids, HPC and SCMC where widely explored for the development of formulation having mucoadhesive properties. since then the use of acrylate polymers for the developments of the mucoadhesion formulation has increased many folds [6].

Anatomy & Physiology of Oral Mucosa

Oral mucosal locale is adhesive in nature and goes about as a lubricant, which is permitting the cells to move comparative with each other with less grating. There are four sites are as follows:

- 1) Buccal cavity
- 2) The sublingual area
- 3) The palate
- 4) Gingival region

It's utilized for drug organization. The utilized site for drug organization of the four-locale referenced over that is the buccal cavity. The anatomic site for drug organization between the cheek and gingival is known as the buccal mucosa. The oral cavity is made out of three layers. The primary layer is the delineated squamous Epithelium, under this layer is basement membrane film. The storm cellular layer overlies the lamina propriety and submucosa. The constitution of the epithelium inside the various locales of the oral cavity show divergence. The epithelium in the slot sense of taste, buccal and sublingual region isn't keratinized, subsequently not containing ceramides and acyl ceramides which are related with giving a boundary work. The mucosa of the buccal & sublingual locale has just modest quantities of ceramide and subsequently more porous. When contrasted with different locales of the oral cavity. A layer of bodily fluid is available on the outer layer of the cells. This assumes a significant part in cell to cell attachment, oral grease just as mucoadhesion of mucoadhesive drug delivery frameworks. The buccal region has a field of smooth and somewhat stable surface, which is appropriate for arrangement of a retentive framework. For buccal drug delivery, grip to the oral mucosa licenses not just the closeness of contact and the chance of further developed drug retention yet in addition the capacity of accomplish an ideal home time at the site of organization These qualities make the buccal mucosa as a more appropriate site for delayed foundational delivery of drugs (Hani M., *et al* 2014).




Figure 1: Overview of oral mucosa

Components and Structural Features of oral cavity

The oral mucosa is for the most part emitted by different organs of oral cavity that are sublingual organ, parotid organ and other salivary organs. The mucus is a clear gel discharged by goblet cell or by uncommon exocrine organs with the mucus cells. The components are given in figure 2:




Figure 2: Composition of Mucus layer

Components	Percentage
Water	95
Glycoproteins & lipids	0.5-5
Mineral salts	1
Free proteins	0.5-1

Table 1. Components of Mucus Layer

Mucus glycoproteins are the high molecular proteins that contain attached oligo- polysaccharide units. The mucus contains following oligosaccharide units:

1. L- fructose
2. D-galactose
3. N-acetyl –D-galactose
4. Sialic acid

Functions of mucus:

1. Protective
2. Lubrication
3. Bio adhesion
4. Cell –cell adhesion
5. Barrier [5]

Mechanism of mucoadhesion

The two material in which one may be artificial such as mucoadhesive polymer and other may be mucin layer of the mucosal tissue are held together by means of interfacial force of attraction is known as mucoadhesion.

Mucoadhesive means artificial substance that is capable of interacting with mucus membrane and being retain on them or holding them together for extended or prolonged time. During the process of adhesion there have two stages identified are given below.

1. Contact Stage
2. Consolidation Stage




Figure 3: Mechanism of mucoadhesion

1) Contact Stage

During at this stage when the mucoadhesive material comes in contact with mucus membrane an intimate wetting occurs between mucoadhesive and mucus membrane. This wetting of mucoadhesive is done by the mucus present in mucosal membrane.

2) Consolidation Stage

By means of different physiochemical forces of attraction such as Vander Waals forces, electrostatic forces and hydrogen bonding. This forces present in mucoadhesive material gets join to the mucus membrane and resulting in long lasting mucoadhesion. This

stage is called as consolidation stage. After these two stages the process of mucoadhesion completes [6].

5.0 Theories of Mucoadhesion



Flow Chart 1: Theories of mucoadhesion

Electronic theory:

According to this theory electron transfer occurs upon contact of an adhesive polymer and the mucus glycoprotein network because of differences in electronic structure. This is proposed to result in the formation of an electronic double layer at the interface, with subsequent adhesion due to attractive forces across the double layer.




Figure 4: Electronic theory of mucoadhesion

Wetting Theory:

It is basically applied to liquid frameworks and things about surface and interfacial energies. It includes the capacity of a fluid to spread suddenly onto a surface as an essential for improvement of bond. The Affinity of liquid for a surface can be found using techniques such as contact angle geometry to measure the contact angle of the liquid on surface, with the general rule being that the lower the contact angle the greater the affinity of liquid to the solid.




Figure 5: Wetting theory of mucoadhesion

Adsorption theory:

According to this theory the attachment of adhesive on the basis of hydrogen bonding and Vander Waals forces. Two types of chemical bonds such as primary covalent & secondary chemical bonds (including electrostatic forces, Vander Waals forces & hydrophobic bonds).

Diffusion theory:

According to this theory the polymer chain and mucus mix to adequate depth to create semi permanent adhesive bond. This process is driven by concentration gradient and is affected by available molecular chain lengths and their mobility's. It depends on the value of molecular weight between cross links and decrease significantly as the cross linking density decreases.




Figure 6: Theory of Diffusion theory

Mechanical theory:

This theory accepts that attachment emergency from an interlocking of fluid cement into anomalies on an unpleasant surface. However, rough surface also provides on increase surface area available for interaction along with an enhanced viscoelastic and plastic dissipation of energy during joint failure, which are thought to be more important in the adhesion process than a mechanical effect.

Fracture theory:

According to this theory of adhesion is related to separation of two surface after adhesion. The fracture strength is equal to adhesive strength, it is given by,

$$G = (E\varepsilon/L)^{1/2} \dots (\text{equation1})$$

Where,

E= Young's module of elasticity

ε = Fracture energy

L = Critical crack length when two surfaces are separated. [2]




Figure 7: Fracture theory of mucoadhesion

Factors affecting mucoadhesion

1) **Molecular weight** Be mucoadhesion strength of a mucoadhesive polymer mainly depend on its molecular weight and polymeric linearity. Generally, for the linear polymers (e.g., Polyethylene glycol), the bio adhesive property are straight away proportional to the molecular weight i.e., PEG-200000 having greater mucoadhesive strength than that of PEG-20000. But in case of nonlinear polymer, the mucoadhesive strength of polymer also can moreover or may not be installed of its molecular weight. Is is mainly because of the reality the helical or coiled structures of such polymer also can moreover protect some of the adhesive group, which can be mainly liable for the adhesive property.

2) **Concentration of polymer** Be interest of a mucoadhesive polymer is a great sized issue of identifying its mucoadhesive electricity. Here is a maximum beneficial interest for a mucoadhesive polymer in which it produces the maximum mucoadhesion. For a few fairly targeted polymeric systems, beyond the maximum beneficial diploma of polymer, the mucoadhesive electricity of polymer starts off advanced to give way notably because of the reality the eye of polymer molecules starts of advanced developing over the molecular interest of the liquid medium in order that there's no similarly chain formation amongst liquid medium and polymer. As an quit end result of this, the polymer particles remain separated from liquid medium, due to this the mucoadhesive electricity of that polymer starts off advanced fallen down.

3) **Flexibility of polymer chains** A more the strength of the mucoadhesive chain motives the more diffusion into the mucus network of buccal cavity. Its effects in stepped forward mucoadhesion. The flexibility of polymer chain decreases with growth in the size of polymer. For an powerful bio adhesion, the polymer chain want to selectively problem into the mucus layer.

4) **Spatial confirmation** The mucoadhesive electricity of a polymer is moreover relying at the conformation or spatial affiliation of polymers i.e., helical or linear. The polymers showing linear conformation having the greater mucoadhesive electricity as have a look at to the polymers showing helical conformation. Because, the helical conformation of polymer also can moreover guard numerous lively groups, which might be at the entire answerable for mucoadhesion, for this reason reducing the mucoadhesive electricity of the polymer.

5) **Swelling or hydration** The proper hydration of mucoadhesive polymer is essential for the popular mucoadhesive strength. With boom in hydration the pore duration of polymer will growth which consequences triggered mobility and advanced interpenetration.

6) **Hydrogen bonding capacity** Hydrogen bonding is every other vital problem for mucoadhesion of a polymer. For mucoadhesion to occur, desired polymers need to have sensible groups which is probably capon a function to form hydrogen bonds. Ability to form hydrogen bonds is due to the presence of (COOH, OH etc.,). Flexibility of the polymer is vital to decorate its hydrogen bonding potential. Polymers which incorporates polyvinyl alcohol, hydroxylase methacrylate and poly (meth acrylic acid) in addition to all their co-polymers are having well hydrogen bonding capacity.

7) **Cross linking density** The cross linking density of the polymer determines its higher molecular weight. The cross linking density indicates the extensive sort of not unusual place molecular weight of the cross linked polymer, which determines the not unusual place pore period. When the cross linking density of polymer is higher, it reduces the pore period of polymer chain which ends up in reduced diffusion of water into the polymer network. It reduced effects with within the decreased penetration of polymer into the mucin and finally decreases the mucoadhesive strength.

8) **Charge** The bio adhesive property of ionic polymer is continuously higher than the of non-ionic polymer. In independent or slightly alkaline medium, the cationic polymer indicates superior mucoadhesive property. It has been hooked up that, cationic immoderate molecular weight polymer which incorporates chitosan very own well bio adhesive [5].

Environment related factors

- **pH of polymer-substrate interface:** The pH of polymer-mucin interface have to be identical as it's miles possible, because, the difference in pH among the 2 structures might also additionally effects withinside the switch of price because of the better pH gradient. It might also additionally act the mucoadhesion.
- **Applied strength:** While setting a buccal mucoadhesive drug transport system, enough electricity have to be implemented if you want to offer an awesome bio adhesive property. Even al even though there aren't any appealing forces among polymer and mucus, then software of excessive stress for sunfficient long term make the polymer come to be bio adhesive with mucus.
- **Initial contact time:** Greater the preliminary touch time among the mucoadhesive polymer and the mucus layer consequences withinside the accelerated swelling in addition to interpenetration of the mucoadhesive polymer chain. Hence, will increase the mucoadhesion electricity of the polymer chain.
- **Moistening:** Moistening is needed to permit the mucoadhesive polymer to unfold over the surface. It creates a community of polymer chains of enough pore size. Through those pores, the interpenetration of polymer and mucin molecules takes

location that outcomes in growing the mobility of polymer chains for the right diffusion of mucoadhesive polymer in mucin layer.

Physiological factors

- **Mucin turnover:** High mucin turnover isn't useful for the mucoadhesive belongings due to following reasons: Be excessive mucin flip over limits the house time of bio adhesive polymer because it detaches from the mucin layer, although it has a very good bio adhesive belongings. High mucin flip over can also additionally produce soluble mucin molecule, for this reason molecule engage with the polymer earlier than they have interaction with mucin layer. Hence there'll now no longer be enough mucoadhesion.
- **Disease state:** In a few sickness states, the secretion of mucus from the mucus membrane receives reduced (e.g., in Dry Mouth Syndrome and in antique age). So that there isn't always sunfficient quantity of mucus gift on the web website online of attachment of mucoadhesive dosage form. Is may also results in mistaken moistening and swelling of polymer. Due to which there may be reduced mucoadhesive power of mucoadhesive dosage form.
- **Rate of renewal of mucosal cells:** Rate of renewal of mucosal cells varies drastically from unique sorts of mucosa. It limits the endurance of bio adhesive structures on mucosal surfaces.
- **Concomitant diseases:** Concomitant sicknesses can regulate the physicochemical homes of mucous or its quantity (for example, hypo and hyper secretion of gastric juice), will increase in frame temperature, ulcer disease, colitis, tissue fibrosis allergic rhinitis, bacterial or fungal contamination and inflammation.
- **Tissue movement:** Tissue motion takes place on intake of liquid and food, speaking, peristalsis withinside the GIT and it impacts the mucoadhesive device mainly in case of gastro retentive dosage forms [5]

Mucoadhesive Dosage Forms

- **Tablet:** Tablets are small, flat, and oval, with a diameter of about 5-eight mm. Unlike the traditional pills, mucoadhesive pills permit for consuming and talking with out principal discomfort. They soften, adhere to the mucosa, and are retained in function till dissolution and or launch is complete. Mucoadhesive pills, in general, have the capacity for use for managed launch drug delivery, however coupling of mucoadhesive houses to pill has extra advantages, for example, it gives green absorption and improved bioavailability of the medicine due a excessive floor to extent ratio and allows a miles greater intimate touch with the mucus layer. Mucoadhesive pills may be tailor-made to stick to any mucosal tissue along with the ones discovered in stomach, as a result providing the opportunities of localized in addition to systemic managed launch of drugs.
- **Films:** Mucoadhesive movies can be desired over adhesive capsules in phrases of flexibleness and comfort. In addition, they are able to stay away from the exceptionally brief house time of oral gels at the mucosa, which can be without problems washed away and eliminated with the aid of using saliva. Moreover, withinside the case of neighborhood transport for oral diseases, the movies additionally assist guard the wound surface, accordingly supporting to lessen pain, and deal with the sickness extra effectively. An best movie need to be flexible, elastic, and soft, but correctly sturdy to resist breakage because of strain from mouth movements. It need to additionally own precise mucoadhesive energy if you want to be retained withinside the mouth for the preferred length of action. Swelling of movie, if it occurs, need to now no longer be too enormous if you want to save you discomfort.
- **Patches:** Patches are laminates which include an impermeable backing layer, a drug-containing reservoir layer from which

the drug is launched in a managed manner, and a mucoadhesive floor for mucosal attachment. Patch structures are much like the ones utilized in transdermal drug delivery. Two strategies used to put together adhesive patches consist of solvent casting and direct milling. In the solvent casting method, the intermediate sheet from which patches are punched is ready via way of means of casting the answer of the drug and polymer(s) onto a backing layer sheet, and in the end permitting the solvent(s) to evaporate. In the direct milling method, system components are homogeneously combined and compressed to the favored thickness, and patches of predetermined length and form are then reduce or punched out. An impermeable backing layer can also be implemented to govern the course of drug release, save you drug loss, and limit deformation and disintegration of the tool in the course of the utility period.

- Gels and ointments:** Semisolid dosage paperwork, inclusive of gels and ointments, have the benefit of clean dispersion at some stage in the oral mucosa. However, drug dosing from semisolid dosage paperwork won't be as correct as from tablets, patches, or films. Poor retention of the gels on the web website online has been conquer with the aid of using the use of mucoadhesive formulations. Certain mucoadhesive polymers, for example, sodium carboxymethylcellulose, carbopol, hyaluronic acid and xanthan gum, go through a segment alternate from liquid to semisolid. This alternate complements the viscosity, which leads to sustained and managed launch of drugs. Hydrogels also are a promising dosage shape for buccal drug shipping. They are fashioned from polymers which might be hydrated in an aqueous surroundings and bodily entrap drug molecules for next gradual launch with the aid of using diffusion or erosion. The software of mucoadhesive gels gives an prolonged retention time withinside the oral cavity, ok drug penetration, in addition to excessive efficacy and affected person acceptability. A important software of adhesive gels is the nearby shipping of medicinal retailers for the remedy of periodontitis, that is an inflammatory and infectious disorder that reasons formation of wallet among the gum and the tooth, and might subsequently motive lack of teeth. It has been advised that mucoadhesive polymers is probably beneficial for periodontitis remedy whilst integrated in antimicrobial-containing formulations which might be effortlessly added into the periodontal pocket with a syringe. HPMC has been used as an adhesive ointment components. Additionally, a distinctly viscous gel become advanced from carpal and hydroxypropyl cellulose for ointment dosage paperwork that might be maintained at the tissue for up to eight hours. (Bind M, et al., 2019).

Marketed Products

Brand name	Active ingredient	Bioadhesive polymer	Dosage form	Company
Aptach	Triamcinolone acetonide	HPC, PAA	Tablet	Teijin Ltd
Buccastem	Prochlorperazine	Xanthan gum, Povidone, Locust bean gum	Tablet	Reckitt Benkiser Plc
Oralin-Generex	Insulin	Unknown	solution	Generex Biotechnology (Phase III trials)
Lauriad	Miconazole	Unknown	Tablet	BioAlliance Pharma (Phase III trials)
Striant SR	Testosterone	Carbomer 934P, Hypromellose, PC	Tablet	Ardana Bioscience Ltd
Suscard	Glyceril trinitrate	Hypromellose	Tablet	Forest Laboratories

Table 2: Marketed Product of mucoadhesion drug delivery system

Conclusion

This review provides the brief knowledge about the oral mucosal drug delivery by discussing briefly the structural feature of mucosa, mechanism of mucoadhesion, various theories of mucoadhesion, general consideration in design of mucoadhesive buccal

dosage forms, permeation enhancers and the various evaluation methods along with the literature Survey of the buccal mucoadhesive drug delivery System.

References

1. S Lairize Mortazavi, John D (1994) Smart; An in vitro method for assessing the duration of Mucoadhesion; J. Control. Release 207-12.
2. NG. Raghvendra Rao, B. Shraavana, Mette Shrikanth Reddy (2013) Overview on Buccal Drug Delivery Systems; J. Pharm. Sic. Res 80-8.
3. P Subramanian (2021) Mucoadhesive Delivery System: A Smart Way to Improve Bioavailability of Nutraceutica 2-22.
4. AB. Bahrain, AK. Khadija (2020) Mucoadhesive Buccal Drug Delivery system: A Review; Am. J. Pharm. Res 274-84.
5. R Singh, D Sharma and Garg (2017) Review on Mucoadhesive Drug Delivery System with Special Emphasis on Buccal Route: An Important Tool in Designing of Novel Controlled Drug Delivery System for the Effective Delivery of Pharmaceutical; J. Dev.Drugs volume 6. Issue 1. 3-12
6. P Mahajan, A Kaur, G Agarwal, SL. Hari Kumar (2013) Mucoadhesive drug delivery system: A Review; Into. Drug. Devers 11-20.
7. Y Akiyama, N Niagara, T kashihara, S Hirai, H Touchy (1995) In vitro and in vivo evaluation of mucoadhesive microspheres

prepared for the Gastrointestinal Tract using polyglycerol esters of fatty acids and a poly (acrylic acid) derivative; Pharm. Res 397-404.

8. Visible A, shade G, Gadge U, Mahajan (2019) An overview on natural polymer based mucoadhesive buccal films for controlled drug delivery; Int. J. Drug. Devers. 1-7

9. . J Gopal Maher, M Tara, P Narayan, Yadav A, Patanaik P, Mishra Singh Yadav (2013) Development and characterization of cellulose- polymethacrylate mucoadhesive film for buccal delivery of carvedilol; Carbohydras. Polyp 172-80.

10. M Montenegro Nicolai, JO Morales (2016) Overview and Future Potential of Buccal Mucoadhesive Films as Drug Delivery System for Biologics; AAPS. Pharm. Sic. Tech 1-8.

11. Shaikh, Thakur Raghu Raj Singh, Mamas Garland A, David Wolfson, Ryan F (2011) Donnelly Mucoadhesive drug delivery systems; J.Pharm. 89-100.

12. Hiba Abdi El Azim, Noah Najee, Ayla Ramadan, Nasal Khalafallah (2015) Liposomal buccal mucoadhesive film for improved delivery and permeation of water- soluble vitamin; Into.J. Pharm. 78-85.

13. S Mansur, P Kesharwani, K Jain, Rakesh K. Tejada N, Karin (2016) Mucoadhesion: A promising approach in drug delivery system; React. Funct. Polyp 151-72.

14. Jain NK (1997) Controlled and Novel Drug Delivery; CBS Publishers. Dist 52-81.

15. J Zhang (1994) An in Vivo Dog Model for Studying Recovery Kinetics of the Buccal Mucosa Permeation Barrier after Exposure to Permeation Enhancers Apparent Evidence of Effective Enhancement without Tissue Damage; Int. J. Pharm 15-22.

16. Smart JD (2005) The basics and underlying mechanisms of mucoadhesion; Adv. Drug. Deli. Rev 1556-68.

17. Hagerstown H, Desman K, Strummed M (2003) Low-frequency dielectric spectroscopy as a tool for studying the compatibility between pharmaceutical gels and mucus tissue; J. Pharm. Sic 92.

18. Lehr CM (1999) Bio adhesive drug delivery Systems-Fundamentals, Novel Approaches and Development; New York: Marcel Dekker p. 1-85.

19. Ahuja A, Khari RK, Ali J (1997) Mucoadhesive drug delivery systems. Drug Dev in Pharm 489-515.

20. Duggan S, Cummins W, Donovan O, Hughes H (2017) Thiolate polymers as mucoadhesive drug delivery system; Ear. J. Pharm. Sic 2-19

21. Gandhi SD, Pandya PR, Mubarak R, Tambuwal T, Shah MA (2011) Mucoadhesive drug delivery systems-an unusual maneuver for site specific drug delivery system. Into J Pharma Sic 132-52.

22. Chowdary KPR, Srinivas L (2000) Mucoadhesive Drug Delivery System: A Review of Current Status. Indian Drugs 400-05.

23. Smart JD The basics and underlying mechanisms of mucoadhesion, Adv. Drug. Deli. Rev 1556-68.

24. Akhtar MH, Gupta J, Mohuddin M., Faisal MD (2015) A comprehensive Review on Buccal Drug Delivery System; Int. J. Of Pharm. Res. and D ev 59-77.

Submit your next manuscript to Annex Publishers and benefit from:

- ▶ Easy online submission process
- ▶ Rapid peer review process
- ▶ Online article availability soon after acceptance for Publication
- ▶ Open access: articles available free online
- ▶ More accessibility of the articles to the readers/researchers within the field
- ▶ Better discount on subsequent article submission

Submit your manuscript at

<http://www.annexpublishers.com/paper-submission.php>