

Rumen Methanogens Community as Drivers of Methane Emission

Teklebrhan T^{*1,2,3}, Tan ZL¹, Wang M¹ and Wang R¹

¹Key Laboratory for Agro-Ecological Processes in Subtropical Region, National Engineering Laboratory for Pollution Control and Waste Utilization in Livestock and Poultry Production, South- Central Experimental Station of Animal Nutrition and Feed Science in the Ministry of Agriculture, Institute of Subtropical Agriculture, The Chinese Academy of Sciences, Changsha, Hunan, P. R. China

²University of Chinese Academy of Sciences, Beijing, China

³Haramaya University School of Animal and Range Sciences, Dire Dawa, Ethiopia

***Corresponding author:** Teklebrhan T, Haramaya University School of Animal and Range Sciences, Dire Dawa, Ethiopia and Key Laboratory for Agro-Ecological Processes in Subtropical Region, National Engineering Laboratory for Pollution Control and Waste Utilization in Livestock and Poultry Production, South- Central Experimental Station of Animal Nutrition and Feed Science in the Ministry of Agriculture, Institute of Subtropical Agriculture, The Chinese Academy of Sciences, Changsha, Hunan, P. R. China, Tel: 15652632823, E-mail: ttmany06@gmail.com

Citation: Teklebrhan T, Tan ZL, Wang M, Wang R (2018) Rumen Methanogens Community as Drivers of Methane Emission. *J Vet Sci Ani Husb* 6(4): 406

Received Date: June 26, 2018 **Accepted Date:** October 24, 2018 **Published Date:** October 26, 2018

Abstract

Among the vast and diverse ruminal micro-biota, Archaea account up to 3-4% of the entire microbial population in rumen of ruminants. *Methanobrevibacter gottschalkii*, *Methanobrevibacter thaueri*, *Methanobrevibacter smithii* and *Methanosphaera stadtmanae* are common ruminal methanogens. Majority of archaea are hydrogenotrophic over acetoclastic methanogen in rumen. Methanogens are endowed with co-enzymes that enable them to produce biogenic methane than other microbes. Biogenic methane produced from ruminants has a significant contribution to global warming. So this review enlightened the complete pictures of methanogens in ruminants and their relationships with other ruminal microbes. The study concluded that the methanogen composition remained variable among ruminant species and these dynamics emanated from animal factors, management and geographical variations.

Keywords: Composition; Diversity; Methanogens; Ruminants; Substrate

Introduction

The inherent characteristics of rumen makes it unique anaerobic habitat to accommodate a community of archaea. Archaea account up to 3-4% of the entire microbial population in the rumen of ruminants (Sharp *et al.* 1998; Yanagita *et al.* 2000 and Ziemer *et al.* 2000) [1-3]. The diversity and population of archaea is very low when compared with bacteria (Lin *et al.* 1997 and Mackie *et al.* 2000) [4,5]. Nearly all of these domain are methanogens, most of which are hydrogenotrophic rather than acetoclastic.

Most of methanogens live freely in rumen liquid or as members of the biofilm adhering to feed particles (Wang *et al.* 2011; Belanche *et al.* 2014 and Jiao *et al.* 2015) [6-8], whereas a small portion of ruminal methanogens are symbionts with protozoa, either ectosymbionts or endosymbionts (Ohene-Adjei *et al.* 2007; Tymensen and McAllister 2012 and Valle *et al.* 2015) [9-11]. There are several possible substrates for methanogenesis in the rumen that originate from bacterial, fungal and protozoa fermentation, including hydrogen (H₂), carbon-dioxide (CO₂), formate, acetate and methyl compounds (Hook *et al.* 2010) [12]. Among which, H₂ and CO₂ are predominant substrates for methanogenesis in the rumen (Hungate *et al.* 1970; Hobson *et al.* 1997; Cavicchioli 2011) [13-15]. There is a strong connection between microbial fermentation processes for H₂ and CH₄ formation by methanogenic archaea in the rumen (Janssen 2010) [16]. Accordingly, methanogens interact with other ruminal microbes, including protozoa (Vogels *et al.* 1980) [17], bacteria (Hegarty and Klieve 1999) [18] and fungi (Brul and Stumm 1994) [19] through interspecies H₂ transfer. Therefore, this review is aimed at understanding of ruminal methanogen community structure, forms, main substrate used and methanogen pathways and hypothesizing a scope for future studies.

Structure and biochemistry of methanogens

Methanogens are a domain of Archaea which has been defined as a separate domain from bacteria and eukaryotes based on the specific 16S ribosomal sequences (Woese *et al.* 1978 and Schafer *et al.* 1999) [20,21]. Methanogens differ from bacteria by the presence of membrane lipids consisting of diether or tetraether linked isoprenoids (De Rosa and Gambacorta 1988) [22] and lack of muramic acid containing peptidoglycans, though some species may have a pseudomuramic acid (Kandler and Hippe 1977) [23]. Another variation between methanogens and bacteria is the structure of archaeal ribosomes. The subunit structure of archaeal ribosomes has a closer resemblance to eukaryotes than true bacteria and the transcription machinery, and the structure of DNA dependent RNA polymerase is also different (Schafer *et al.* 1999) [21].

Forms of methanogens

Free living methanogens

Most of methanogens are free living which is not associated with ruminal protozoa or fungi (Belanche *et al.* 2014) [7]. Majority of free-living methanogens are integrated into biofilm on the surface of feed particles where H₂-producing bacteria actively produce H₂ (Leng 2014) [24]. These methanogens may not be inhibited as much as free-living peers by anti-methanogenic inhibitors. None of the Sequences related to *Methanobacterium* and *Methanosphaera* have been recovered from rumen protozoa and, thought to be free-living methanogens. In contrast, a significant portion (32.8%) of the *Methanobrevibacter* sequences archived in Ribosomal Database Project (RDP) was recovered from protozoa. However, *Methanobrevibacter* accounts for at least 65% of the rumen methanogens in which significant portion of the *Methanobrevibacter* sequences are not free living may simply reflect the probability of sequence recovery, rather than a selective association between rumen protozoa and *Methanobrevibacter*. It is reflected by smaller number of 16S rRNA gene sequences recovered from protozoa than from rumen content or fluid. The result could also arise from difficulties associated with obtaining archaeal DNA from protozoal cells (Belanche *et al.* 2014) [7].

Protozoa associated methanogens (PAM)

Most species of rumen ciliate protozoa contain hydrogenosomes, a unique type of membrane bounded organelles producing H₂ by malate oxidization (Muller 1993) [25]. These organelles can attract some methanogens as endosymbionts (Janssen and Kirs 2008) [26]. Hydrogen generated by rumen protozoa could be utilized by PAM, which benefits both parties. Methanogens have been found internally (Finlay *et al.* 1994) [27] and externally in association with protozoa (Vogels *et al.* 1980 and Krumholz *et al.* 1983) [17,28]. Stumm *et al.* (1982) [29], observed that the frequency of ectosymbiotic methanogens was affected by relative contribution of H₂ production by rumen ciliates and H₂-producing bacteria. Based on fluorescence in situ hybridization analysis, about 16% of the rumen ciliates contained methanogens inside their cells (Lloyd *et al.* 1996) [30]. A possible explanation for low incidence is that intracellular association may be transient rather than permanent. However, earlier studies indicated that rumen ciliates do not have endosymbiotic methanogens though they might have ectosymbiotic methanogens (Williams and Coleman, 1992) [31]. The difficulty in distinguishing engulfed methanogens from true endosymbiotic methanogens presents a challenge in determining if rumen ciliates possess true endosymbiotic methanogens and or bacteria. Because of labor intensive procedures, PAM are mostly identified using DNA based methods, and only one strain of methanogens (isolates MB-9, related to *Methanobrevibacter ruminantium*) has been reported to be associated with a ciliate in the rumen of sheep (Tokura 1999) [32]. Among the methanogen sequences of rumen origin archived in the RDP database (Release 11, Update 3), only a very small proportion (5.3%) was recovered from washed protozoa cell (Chagan *et al.* 1999; Tokura *et al.* 1999; Regensbogenova *et al.* 2004; Ohene-Adjei *et al.* 2004; Tymensen and McAllister 2012) [10,32-35]. *Methanobrevibacter* and *Methanomicrobium* were the first and second largest genera reported to be PAM, and they accounted for 32.8% and 23.0% of the total sequences, respectively. *Methanomicrobium* is better represented in PAM sequences than total archaeal sequences (7.7%), species of both taxa may be among the predominant PAM. However, Janssen and Kirs (2008) [26], declared that the above results may be biased because only a small number of PAM sequences were obtained from selected protozoa. Besides, the PAM sequences may be contaminated with sequences of non-PAM.

Rumen methanogens population

Most of the ruminant methanogens are known to be resident of rumen and belong to three principal genera including *Methanobrevibacter*, *Methanomicrobium* and 'rumen cluster C' (Paul *et al.* 2012) [36], while the rest belong to minority genera such as *Methanimicrococcus*, *Methanosarcina* and *Methanobacterium* (Janssen and Kirs 2008; St-Pierre and Wright 2013) [26,37]. A varieties of methanogens can be found in the rumen (Janssen and Kirs 2008; Gemma *et al.* 2015) [26,38]. Gemma *et al.* (2015) noted that 77.7% of methanogens in the rumen were hydrogenotrophic, while 22.1% had an ability to grow with H₂ and methyl groups derived from methanol or methylamines. According to earlier study by Kim *et al.* (2011) [39], the Archaea in rumen of domesticated livestock deposited in the Ribosomal Database Project (RDP) accounts 3516 archaeal sequences. During that time, the species of archaea in the rumen is estimated to be approximately 949 and with estimated maximum number of species 1469 (Kim *et al.* 2011) [39]. Nearly all the archaeal sequences were assigned to the phylum Euryarchaeota which was represented by about 670 genus-level OTUs and 1000 species-level OTUs. Compared with other anaerobic habitats, greater than

100 species of methanogens of 28 genera have been isolated, the diversity and species richness of ruminal methanogens are quite low, reflecting the highly selective ruminal environment for methanogens. Rarefaction analysis of the sequences in the database indicated that the coverage of the diversity at species level was 65% for archaea in 2011. However, very recently ruminal methanogens accounts about 8623 archaeal sequences of rumen origin (RDP Release 11, update 3). About 90% of these sequences were assigned to methanogens (RDP Release 11, update 3). These sequences are classified onto ten known genera, majority of sequences were under *Methanobrevibacter* followed by *Methanosphaera*, *Methanomicrobium* and *Methanobacterium*. The order Thermoplasmatales, which was previously referred to as the rumen cluster C (RCC) group, is represented by few sequence of the total archaeal sequences.

Among the sequences in the data bases, eight species of methanogens have been isolated into pure cultures of *Methanobacterium formicicum*, *Methanobacterium bryantii*, *Methanobrevibacter ruminantium*, *Methanobrevibacter millerae*, *Methanobrevibacter olleyae*, *Methanomicrobium mobile*, *Methanoculleus olentangyi*, and *Methanosarcina barkeri* (McAllister *et al.*, 1996; Janssen and Kirs, 2008; Zhou *et al.*, 2008) [26,40,41]. Moreover, in 2013 five new species were isolated, including *Methanobrevibacter boviskoreani* (Korean native cattle) (Lee *et al.* 2013) [42], *Methanobacterium beijingense* (goat), *Methanoculleus marisnigri* (Indian crossbred cattle), *Methanoculleus bourgensis* (Holstein cattle), and *Methanosarcina mazei* (isolated from the rumen of Korean Hanwoo cattle) (RDP database). Besides, one Thermoplasmatales-like pyrrolysine-dependent methanogen was isolated from bovine (GenBank access number: CP002916). Collectively, 16S rRNA gene sequences from cultured methanogens only accounted for 0.7% of the total archaeal sequences of rumen origin. Most of the isolates are members of the family Methanobacteriaceae and several taxa do not have a single cultured representative. Therefore, there are many uncultured species/ strains of methanogens have been isolated with culture independent techniques like 16S rRNA gene clone libraries and DNA sequence analysis (Klieve *et al.* 2009; Ouwerkerk *et al.* 2008; Wright *et al.* 2008). The possible reason might be these species owing to very specific requirements for particular substrates and other physiological conditions.

Genera and species

The majority (92.3%) of archaea are classified into three genera such as *Methanobrevibacter* (61.6%), *Methanomicrobium* (14.9%), as well as a *Methanomassiliicoccales* (15.8%) (Sirohi *et al.* 2010) [43]. Among which *Methanobrevibacter* was the dominant genera of methanogens followed by *Methanosphaera* and *Methanobacterium* in ruminants (Wright *et al.*, 2004, 2008; Nicholson *et al.*, 2007; Ouwerkerk *et al.*, 2008; Snelling *et al.*, 2014; St-Pierre *et al.*, 2015; Zhou *et al.* 2017a, 2017b and Wang *et al.* 2017) [44-52]. On the other hand, few reports indicated that the dominant ruminal methanogens belonged to rumen cluster C in dairy cows accounting 71% of the total sequences (Wang *et al.* 2016) [53]. Similarly, in Tibetan ruminants appears to affiliate with rumen cluster C or recently *Methanoplasmatales* (Iino *et al.*, 2013 and Huang *et al.*, 2016) [54,55]. This review suggests that there is high range of variation with regards to methanogen species among ruminant species and this attributes to type of feed, species, location of animals, type of primers, isolation techniques used by different scientist are highly variable. Generally, this review summarized that the relative abundance of the methanogens genera and respective species in ruminant are shown in order of importance in Table 1.

Item	References
<i>Methanobrevibacter</i> genera <i>Methanobrevibacter ruminantium</i>	Tatsuoka <i>et al.</i> 2004; Wright <i>et al.</i> 2007; Nicholson <i>et al.</i> 2007; wang <i>et al.</i> 2017
<i>Methanobrevibacter smithii</i> , <i>methanobrevibacter gottschalkii</i> , <i>Methanobrevibacter millerae</i> or <i>ethanobrevibacter thaueri</i>	Zhou <i>et al.</i> 2009; St-Pierre <i>et al.</i> 2015; Henderson <i>et al.</i> 2015 and Gemma <i>et al.</i> 2015; Huang <i>et al.</i> 2016
<i>Methanosarcina barkeri</i>	McAllister <i>et al.</i> 1996; Moss <i>et al.</i> 2000; Boadi <i>et al.</i> 2004; Ouwerkerk <i>et al.</i> 2008
<i>Methanomicrobium</i> genera <i>Methanomicrobium mobile</i>	Lin <i>et al.</i> 1997; Yanagita <i>et al.</i> 2000; Tajima <i>et al.</i> 2001; Regensbogenova <i>et al.</i> 2004; Shin <i>et al.</i> 2004 and Chaudhary and Sirohi 2008;
Ruminant cluster C (RCC) genera Rumen cluster spp.	Tajima <i>et al.</i> 2001; Wright <i>et al.</i> 2006, 2007; Janssen and Kirs 2008;
<i>Methanomassiliicoccales</i> genera <i>Methanosarcina</i> spp	Lin <i>et al.</i> 1997; Yanagita <i>et al.</i> 2000; Skillman <i>et al.</i> 2004
<i>Methanosphaera stadmanae</i>	Sharp <i>et al.</i> 1998 and Wright <i>et al.</i> 2004

Source: 2018 review

Table 1: List of methanogens genera and their respective species

Substrates for methanogens

Majority of rumen archaea use H₂ and CO₂ as main substrates for formation of methane (CH₄). The broad diversity of methanogens in rumen would suggest factors other than H₂ and CO₂ in determining structure of the community. Hydrogenotrophic methanogens use CO₂ or acetate as their carbon source and H₂ as main electron donor play a dominant role during methanogenesis. While During formatotrophic methanogenesis, four molecules of formate are oxidized to form CO₂ by formate dehydrogenase (FDH). Members of the order Methanosarcinales (*Methanosphaera* spp.) and Methanobacteriales have been classified into methylotrophs that use methyl containing compounds such as methanol, methylamines or dimethylsulfide to produce CH₄. Acetate is preferred

substrate for *Methanosarcina* and *Methanosaeta* that produce CH_4 via acetoclastic pathway (Liu and Whitman 2008) [56]. During acetoclastic methanogenesis, acetate splits to form carboxyl compounds that become oxidized to CO_2 . Methyl groups from CO_2 then enters the hydrogenotrophic pathway to form CH_4 . Contribution of methyl groups and acetate as substrates for methanogenesis in rumen is likely to be minimal as methanogens that depend on these conversions for producing CH_4 have a very slow growth rate in vitro, suggesting short retention time exerted by normal rumen conditions would prevent them from thriving (Janssen and Kirs 2008) [26]. Miller (2001) [57] noted that *Methanospaera spp.*, as well as *Methanospaera stadmanae* from the human colon and *Methanospaera cuniculi* isolated from the contents of rabbit rectum, can utilize only methanol with H_2 as substrate for growth unlike other methanogenic archaea. Because, *Methanospaera stadmanae* genome does not contain molybdopterin biosynthesis genes and therefore, it is unable to synthesize active formylmethanofuran dehydrogenase, which forms formylmethanofuran from CO_2 (Fricke *et al.* 2006) [58]. In addition, this archaea also lacks genes for carbon monoxide dehydrogenase/acetyl coenzyme A synthase complex (Fricke *et al.* 2006) [58], which may explain why *Methanospaera stadmanae* cannot use acetate as a methanogenic substrate. *Methanospaera stadmanae* were identified from captive Sumatran orangutans (*Pongo abelii*) fecal samples. This animal eats fresh fruit as 50% of its daily diet and degradation of fruit pectin could provide methanol as a methanogenic substrate to *Methanospaera stadmanae* to outcompete other methanogenic groups. *Methanospaera stadmanae* can utilize only methanol with H_2 for CH_4 production, while *Methanobrevibacter smithii* not only uses CO_2 , H_2 and/or formate but also contains enzymes facilitating methanol and ethanol utilization (Samuel *et al.* 2007) [59].

Apart from the above common substrates, phosphatidylcholine is also a compound of the plant membrane materials and is rapidly degraded by rumen microorganisms to liberate choline (Neill *et al.* 1978) [60]. Choline is subsequently degraded to trimethylamine. The methyl groups of trimethylamine and methylamine are converted to CH_4 , apart from other *Methanobrevibacter spp.* including *Methanobrevibacter ruminantium*, *Methanosarcina barkeri* utilize trimethylamine, dimethylamine, and methylamine as substrates for methanogenesis (Hippe *et al.* 1979; Patterson and Hespell 1979) [61,62]. Patterson and Hespell (1979) revealed that when other substrates are unavailable in the bovine rumen, the presence of methylamine compounds may allow noncompetitive growth of *Methanosarcina barkeri*.

Methanogenesis pathways

Fermentation of carbohydrates results in accumulation of hydrogen, the removal of which through methanogenesis pathways by methanogens allows microorganisms to degrade substrates continuously and supports the rumen anaerobic fermentation process and this occurs by oxidation of sugars via Embden-Meyerhof-Parnas pathway in bacteria, fungi and protozoa, electron carrying cofactors such as NADH must be reoxidized to NAD^+ to allow fermentation to continue (McAllister and Newbold 2008) [63]. Three types of methanogenic metabolic pathways are involved in CH_4 synthesis, namely hydrogenotrophic (reduction of CO_2 coupled to the oxidation of H_2), methylotrophic (conversion of methyl-group containing compounds) and acetoclastic (Rother and Krzycki 2010; Ferry 2011) [64,65].

Hydrogenotrophic pathway is generally recognized as the main pathway to remove hydrogen, through which methanogens can utilize H_2 as electron donor to reduce CO_2 to CH_4 . Newly recognized methanogens (e.g., rumen cluster C) use a range of methyl donor compounds and CO_2 for CH_4 production, suggesting that other pathways maybe identified (Poulsen *et al.* 2013) [66]. The draft genome of *Candidatus Methanomethylphilus Mx1201*, a methanogen isolated from human gut belonging to the rumen cluster C, more recently categorized into the order Methanomassiliococcales (Dridi *et al.* 2012) [67], contains genes for methylotrophic methanogenesis from methanol and tri-, di- and monomethylamine (Borrel *et al.* 2012) [68]. In artificial systems, such as biogas production facilities, acetate is recognized as an important substrate for methanogens, which refers to as acetoclastic methanogenesis (Weiland 2010) [69]. A comprehensive understanding of the functionality of methanogens and their CH_4 producing pathways may provide insights into effective CH_4 abatement strategies.

Methanogenesis is unique to the methanogens, this ability is due to the presence of three coenzymes (Figure 1), which have not been found in other microorganisms namely, coenzyme M, coenzyme B (Factor B) and CoM-S-S-CoB. All of these coenzymes are involved in various oxidation and reduction processes during CH_4 formation (Baker 1999; Boadi *et al.* 2004) [70,71]. The formation of CH_4 from methanol is similar except that from methanol there is either direct formation of CH_4 through coenzyme M, or by methanol oxidation to CO_2 through a pathway which initially proceeds in the opposite direction to CO_2 reduction pathway, before transfer of the methyl group to coenzyme-M, which is then reduced to CH_4 (Ferry 1993) [72]. Thus, the last step in the formation of CH_4 from all pathways proceeds through coenzyme M which is reduced by the methyl-CoM reductase (mcr) enzyme and a cofactor F430 exclusive to the archaea which acts as a prosthetic group to enzyme mcr (Thauer 1998) [73]. It can thus be concluded that methanogens have highly conserved 16S ribosomal sequences which make them a distinct group from other microorganisms. These unique sequences can thus be targeted to perform culture independent molecular analysis of the methanogens.




Figure 1: Methanogenesis and role of co-enzymes

It summarizes methanogenesis using the exclusive enzymes mainly coenzyme M which is reduced by an enzyme mcr to produce CH₄ production from all substrates

Association of methanogens and microbes

Associations mainly occur via pools of common metabolites, especially where end products of one group form the substrates of another group. Methanogens interact with other ruminal microbes, including protozoa (Vogels *et al.* 1980) [17], bacteria (Hegarty and Klieve 1999) and fungi (Brul and Stumm 1994) [18,19]. Overall, such interaction benefits rumen fermentation to prevent accumulation of reducing equivalents and inhibition of rumen fermentation and facilitate normal fermentation process in rumen (Kong *et al.* 2013) [74]. Methanogens can associate with fungi, but little is known about fungal-associated methanogens. An early study suggests that rumen fungi do not have endosymbiotic methanogens although they may have ectosymbiotic methanogens (Marvin-Sikkema *et al.* 1992) [75]. In a recent study, species of *Methanobrevibacter* were detected in cultures of *Piromyces*, *Anaeromyces*, and *Neo-callimastix* (Jin *et al.* 2014) [76]. In addition, methanogens were evident in some rumen fungal cultures, but it was not reported if methanogens and fungi had any type of physical association (Jin *et al.* 2014) [76]. All rumen fungi contain hydrogenosomes and methanogens use H₂ to produce CH₄ (Orpin 1988) [77], but concrete evidence is needed to determine if rumen fungi carry true endosymbiotic methanogens.

A recent study investigated associations between bacteria, archaea, and protozoa, but no strong correlations between archaea and protozoa (Gemma *et al.* 2015) [38], though in earlier study methanogens are known to colonize protozoa, and this mutualistic relationship is believed to enhance CH₄ formation in the rumen (Newbold *et al.* 1995) [78]. Moreover, the occurrence of specific symbiosis between methanogens and rumen ciliate protozoa has been speculated on, but till further verification is required (Hackstein 2010) [79]. Lack of strong co-occurrence patterns indicate that these important associations are probably non-specific, or occur at a strain level and, further investigation is required to validate those reports, as mechanisms that mediate the colonization of protozoa by archaea remain to be elucidated. These could have interesting evolutionary aspects if they allow non-specific interactions to form or are mediated by strain-specific mechanisms that confer different partner specificities within archaeal or protozoal species. In addition, lack of strong association patterns between protozoa and major methanogen groups on the other way, suggests that conserved mechanisms may mediate the interactions between H₂ producing and H₂ consuming microbes, allowing flexible interactions. This may aid CH₄ mitigation research, since interfering with these potentially universal mechanisms could slow the rate of H₂ transfer and so slow CH₄ formation (Stams and Plugge 2009) [80].

Cellulolytic bacteria also have association with methanogens, Minato *et al.* (1992) [81] hypothesized that cellulolytic bacteria supply H₂ to methanogens. In line to this, Morgavi *et al.* (2010) [82], noted that integration of methanogens into bacterial biofilms on feed particles itself represents a form of interaction, and most fermentative ruminal bacteria produce CO₂ and H₂. Thus, rumen bacteria and methanogens interact mutualistically through interspecies H₂ transfer. Such interspecies H₂ transfer was demonstrated in co-cultures of methanogens with *Ruminococcus albus* (Wolin *et al.* 1997) [83], *R. flavefaciens* (Latham *et al.* 1977) [84] and *Selenomonas ruminantium* (Scheifinger *et al.* 1975) [85]. Consequently, interaction between rumen bacteria and methanogens affects energy conservation, volatile fatty acid profiles, and CH₄ production by rumen microbiome. On the

other hand, no strong associations were found between most abundant bacteria and archaea (Gemma *et al.* 2015) [38], It was surprising findings, since rumen bacteria degrade feed and produce the substrates for methanogens, mainly H₂ and methyl groups. However, there was positive associations between some less abundant bacteria and archaea, the bacteria were succinate-producing Succinivibrionaceae, succinate-using Dialister, and the amino-acid-fermenting Acidaminococcus, and methanogens related with Methanomassiliococcaceae, Methanosphaera spp. A4, and Methanobrevibacter boviskoreani. Succinivibrio spp. degrade pectin (Bryant 1956) [86]. Methanomassiliococcaceae (Paul *et al.* 2012) [87] and Methanosphaera (Miller and Wolin 1985) [88]. Other associations were also between the methylotrophic methanogens Methanosphaera sp. ISO3-F5 and different bacteria, including members of Lachnospiraceae. These associations may be based on the ability of Lachnospiraceae to degrade pectin and so provide methanol as a substrate for methylotrophs (Dehority 1969) [89]. The associations between other Methanomassiliococcaceae groups and various unclassified members of Bacteroidales also suggest the possibility of yet further methanol-dependent metabolic interactions [90-96].

Conclusion

It might be possible to deduce that with the exception of rare studies, the genera Methanobrevibacter and methanobacterium are considered as the dominant methanogenic group followed by RCC in ruminants. However, methanogen community structure is so dynamic depending on animal and other environmental factors. This review finally, hypothesized that most of the previous studies are focused on methanogen community only in rumen of ruminants focusing with obvious pathway. However, little is known about the distribution of methanogens across the gastro-intestinal tracts of ruminates. Hence, to discover some novel genes, future study is warranted to address the issues to have a complete understanding of methanogens in ruminants. Moreover, methanogen community structure and methane production pathway using format as a substrate is remained to be unknown and future works are needed in addressing these gaps.

Acknowledgments

The author is highly grateful for CAS-TWAS scholarship for sponsoring the study in China. I would like also to thank the institute of Subtropical Agriculture, University of Chinese Academy of Sciences for allowing me to have access to different data bases.

References

1. Sharp R, Ziemer CJ, Stern MD, Stahl DA (1998) Taxon-specific associations between protozoal and methanogen populations in the rumen and a model rumen system. *FEMS Microbiol Ecol* 26: 71-78.
2. Ziemer CJ, Sharp R, Stern MD, Cotta MA, Whitehead TR, et al. (2000) Comparison of microbial populations in model and natural rumens using 16S ribosomal RNA-targeted probes. *Environ Microbiol* 2: 632-43.
3. Lin C, Raskin L, Stahl DA (1997) Microbial community structure in gastrointestinal tracts of domestic animals: comparative analyses using rRNA-targeted oligonucleotide probes. *FEMS Microbiol Ecol* 22: 281-94.
4. Mackie RI, Aminov RI, White BA, McSweeney CS (2000) Molecular ecology and diversity in gut microbial ecosystems. Ruminant physiology: digestion, metabolism, growth and reproduction. CAB International, Oxford 61-77.
5. Wang P, Qi M, Barboza P, Leigh MB, Ungerfeld E, et al. (2011) Isolation of high-quality total RNA from rumen anaerobic bacteria and fungi, and subsequent detection of glycoside hydrolases. *Can J Microbiol* 57: 590-8.
6. Belanche A, de la Fuente G, Newbold CJ (2014) Study of methanogen communities associated with different rumen protozoal populations. *FEMS Microbiol Ecol* 90: 663-77.
7. Jiao J, Li X, Beauchemin KA, Tan Z, Tang S, et al. (2015) Rumen development process in goats as affected by supplemental feeding v. grazing: age-related anatomic development, functional achievement and microbial colonisation. *Br. J. Nutr.* 113: 888-900.
8. Ohene-Adjei, Teather SR, Ivan M, Forster M, Postinoculation RJ (2007) Protozoan Establishment and Association Patterns of Methanogenic Archaea in the Ovine Rumen. *Appl Environ Microbiol* 73: 4609-18.
9. Tymensen LD, McAllister TA (2012) Community Structure Analysis of Methanogens Associated with Rumen Protozoa Reveals Bias in Universal Archaeal Primers. *Appl Environ Microbiol* 78: 4051-6.
10. Valle ER, Henderson G, Janssen PH, Cox F, Alexander TW, et al. (2015) Considerations in the use of fluorescence in situ hybridization (FISH) and confocal laser scanning microscopy to characterize rumen methanogens and define their spatial distributions. *Can J Microbiol* 61: 417-28.
11. Hook SE, Wright ADG, McBride BW (2010) Methanogens: methane producers of the rumen and mitigation strategies. *Archaea* P 1 -11.
12. Hungate RE, Smith W, Bauchop T, Yu I, Rabinowitz JC (1970) Formate as an intermediate in the bovine rumen fermentation. *J Bacteriol* 102: 389-97.
13. Hobson PN, Stewart CS (2010) The rumen microbial ecosystem. London: Chapman and Hall, 1997.
14. Janssen PH (2010) Influence of hydrogen on rumen methane formation and fermentation balances through microbial growth kinetics and fermentation thermodynamics. *Anim Feed Sci Technol*. 160: 1-22.
15. Vogels GD, Hoppe WF, Stumm CK (1980) Association of methanogenic bacteria with rumen ciliates. *Appl Environ Microbiol* 40: 608-12.
16. Hegarty R, Klieve A (1999) Opportunities for biological control of ruminal methanogenesis. *Crop Pasture Sci* 50: 1315-20.
17. Brul S, Stumm CK (1994) Symbionts and organelles in anaerobic protozoa and fungi. *Trends Ecol Evol* 9: 319-24.
18. Woese CR, Magrum LJ, Fox GE (1978) Archaeobacteria. *J Mol Evol* 11: 245-52.
19. Schafer G, Engelhard M, Muller V (1999) Bioenergetics of the Archaea. *Microbiology and Molecular Biology Reviews* 63: 570-620.
20. De Rosa M, Gambacorta A (1988) The lipids of archaeobacteria. *Progress in Lipid Research* 27: 153-75.
21. Kandler O, Hippe H (1977) Lack of peptidoglycan in the cell walls of *Methanosarcina barkeri*. *Arch Microbiol* 113: 57-60.

22. Leng R (2014) Interactions between microbial consortia in biofilms: A paradigm shift in rumen microbial ecology and enteric methane mitigation. *Anim Prod Sci* 54: 519-43.
23. Muller M (1993) Review article: The hydrogenosome. *J Gen Microbiol* 139: 2879-89.
24. Janssen PH, Kirs M (2008) Structure of the Archaeal Community of the Rumen. *Appl Environ Microbiol* 74: 3619-5.
25. Finlay BJ, Esteban G, Clarke KJ, Williams AG, Embley TM, et al. (1994) Some rumen ciliates have endosymbiotic methanogens. *FEMS Microbiology Letters* 117: 157-61.
26. Krumholz LR, Forsberg CW, Veira DM (1983) Association of methanogenic bacteria with rumen protozoa. *Canadian J Microbiol* 29: 676-80.
27. Stumm CK, Gijzen HJ, Vogels GD (1982) Association of methanogenic bacteria with ovine rumen ciliates. *Bri J Nutr* 47: 95-9.
28. Tokura M, Chagan I, Ushida K and Kojima Y (1999) Phylogenetic study of methanogens associated with rumen ciliates. *Current Microbiology* 39: 123-8.
29. Chagan I, Tokura M, Jouany JB, Ushida K (1999) Detection of methanogenic archaea associated with rumen ciliate protozoa. *J Gen Appl Microbiol* 45: 305-8.
30. Regensbogenova M, McEwan N, Javorsky P, Kisidayova S, Michalowski T, et al. (2004) A re-appraisal of the diversity of the methanogens associated with the rumen ciliates. *FEMS Microbiol Lett* 238: 307-13.
31. Adjei O, Opoku C (2004) Urinary tract infections in African infants. *Int J Anti Mic Ag* 24: 32-4.
32. Paul K, Nonoh JO, Mikulski L, Brune A (2012) "Methanoplasmatales", thermoplasmatales-related Archaea in termite guts and other environments, are the seventh order of methanogens. *Appl Environ Microbiol* 78: 8245-53.
33. St-Pierre B, Wright ADG (2013) Diversity of gut methanogens in herbivorous animals. *Animal* 7: 49-56.
34. Kim M, Morrison M, Yu Z (2011) Status of the phylogenetic diversity census of ruminal microbiomes. *FEMS Microbiology Ecology* 76: 49-63.
35. McAllister TA, Okine EK, Mathison GW, Cheng KJ (1996) Dietary, environmental and microbiological aspects of methane production in ruminants. *Canadian J Anim Sci* 76: 231-43.
36. Zhou M, Hernandez-Sanabria E, Guan LL (2009) Assessment of the microbial ecology of ruminal methanogens in cattle with different feed efficiencies. *Appl Environ Microbiology* 75: 6524-33.
37. Lee JH, Kumar S, Lee GH, Chang DH, Rhee MS, et al. (2013) *Methanobrevibacter boviskoreani* sp. nov., isolated from the rumen of Korean native cattle. *Intl J Syst Evol Microbiol*. 63: 4196-201.
38. Chaudhary PP, Sirohi SK (2009) Dominance of *Methanomicrobium* phylotype in methanogen population present in Murrah buffaloes (*Bubalus bubalis*). *Let Appl Microbiol* 49: 274-7.
39. Wright ADG, Williams AJ, Winder B, Christophersen CT, Rodgers SL, et al. (2004) Molecular diversity of rumen methanogens from sheep in Western Australia. *Appl Environ Microbiol* 70: 1263-70.
40. Wright ADG, Auckland CH, Lynn DH (2007) Molecular diversity of methanogens in feedlot cattle from Ontario and Prince Edward Island, Canada. *Appl Environ Microbiol* 73: 4206-10.
41. Nicholson MJ, Evans PN, Joblin KN (2007) Analysis of methanogen diversity in the rumen using temporal temperature gradient gel electrophoresis: identification of uncultured methanogens. *Microb Ecol* 54: 141-50.
42. Ouwerkerk D, Turner AF, Klieve AV (2008) Diversity of methanogens in ruminants in Queensland. *Aus J Exp Agri* 48: 722-5.
43. Snelling TJ, Genc B, McKain N, Watson M, Waters SM, et al. (2014) Diversity and community composition of methanogenic archaea in the rumen of Scottish upland sheep assessed by different methods. *PLoS One* 9: e106491.
44. St-Pierre B, Cersosimo LM, Ishaq SL, Wright ADG (2015) Toward the identification of methanogenic archaeal groups as targets of methane mitigation in livestock animals *Frontiers in Microbiology* 6: 776. doi: 10.3389/fmicb.2015.00776.
45. Zhou Z, Meng Q, Li S, Jiang L, Wu H (2017a) Effect of urea-supplemented diets on the ruminal bacterial and archaeal community composition of finishing bulls. *Appl Microbiol Biotechnol* DOI 10.1007/s00253-017-8323-4.
46. Zhou Z, Fang L, Meng Q, Li S, Chai S, et al. (2017 b) Assesment of bacterial and Archeal community structure in Yak (*Bos grunniens*). *Front Microbiol* 8: 179.
47. Wang Z, Elekwachi CO, Jiao J, Wang M, Tang S, et al. (2017) Investigation and manipulation of metabolically active methanogen community composition during rumen development in black goats. *Scientific Reports* 7: 422 doi: 10.1038/s41598-017-00500-5.
48. Wang P, Zhao S, Wang X, Zhang Y, Zheng N, et al. (2016) Ruminal methanogen community in dairy cows fed agricultural residues of corn stover, rapeseed, and cottonseed meals *J. Agric Food Chem* DOI: 10.1021/acs.jafc.6b00708.
49. Iino T, Tamaki H, Tamazawa S, Ueno Y, Ohkuma M, et al. (2013) *Candidatus methanogramma caenicola*: a novel methanogen from the anaerobic digested sludge, and proposal of *Methanomassiliicoccales* fam. nov. and *Methanomassiliicoccales* ord. nov., for a methanogenic lineage of the class 'Thermoplasmata,' *Microbes and Environments* 28: 244-50.
50. Huang XD, Martinez-Fernandez G, Padmanabha J, Long R, Denman SE, et al. (2016) Methanogen Diversity in Indigenous and Introduced Ruminant Species on the Tibetan Plateau. *Archaea* 5916067.
51. Liu Y, Whitman WB (2008) Metabolic, phylogenetic, and ecological diversity of the methanogenic archaea. *Ann N Y Acad Sci* 1125: 171-89.
52. Boone DR, Castenholz RW, Garrity GM (2001) Genus II *Methanobrevibacter* Balch and Wolfe 1981, 216VP. In: *Bergey's manual of systematic bacteriology*, New York, NY: Springer, USA.
53. Fricke WF, Sedorf H, Henne A, Krüer M, Liesegang H, et al. (2006). The genome sequence of *Methanosphaera stadtmanae* reveals why this human intestinal archaeon is restricted to methanol and H₂ for methane formation and ATP synthesis. *J Bacteriol* 188: 642-58.
54. Samuel BS, Hansen EE, Manchester JK, Coutinho PM, Henriessat B, et al. (2007) Genomic and metabolic adaptations of *Methanobrevibacter smithii* to the human gut. *Proc Natl Acad Sci USA* 104: 10643-8.
55. Neill AR, Grime DW, Dawson RM (1978) Conversion of choline methyl groups through trimethylamine into methane in the rumen. *Biochem J* 170: 529-35.
56. Hippe H, Caspari D, Fiebig K, Gottschalk G (1979) Utilization of trimethylamine and other N-methyl compounds for growth and methane formation by *Methanosarcina barkeri*. *Proc Natl Acad Sci USA* 76: 494-8.
57. Patterson JA, Hespell RB (1979) Trimethylamine and methylamine as growth substrates for rumen bacteria and *Methanosarcina barkeri*. *Curr Microbiol* 3: 79-83.
58. McAllister TA, Newhold CJ (2008) Redirecting rumen fermentation to reduce methanogenesis. *Aust J Exp Agric* 48: 7-13.

59. Rother M, Krzycki JA (2010) Selenocysteine, pyrrolysine, and the unique energy metabolism of methanogenic archaea. *Archaea* 45: 36-42.
60. Ferry JG (2011) Fundamentals of methanogenic pathways that are key to the biomethanation of complex biomass. *Curr Opin Biotechnol* 22: 351-7.
61. Poulsen MC, Schwab BB, Jensen RM, Engberg A, Spang N, et al. (2013) Methylophilic methanogenic Thermoplasmata implicated in reduced methane emissions from bovine rumen. *Nat Commun* 4: 14-28.
62. Dridi B, Fardeau M L, Ollivier B, Raoult D, Drancourt M (2012) *Methanomassiliicoccus luminyensis* gen. nov., sp. nov., a methanogenic archaeon isolated from human faeces. *Int J Syst Evol Microbiol* 62: 1902-7.
63. Borrel G, Harris HM, Tottey W, Mihajlovski A, Parisot N, et al. (2012) Genome sequence of "Candidatus Methanomethylophilus alvus" Mx1201, a methanogenic archaeon from the human gut belonging to a seventh order of methanogens. *J Bacteriol* 194: 6944-5.
64. Weiland P (2010) Biogas production: current state and perspectives. *Appl Microbiol Biotechnol* 85: 849-60.
65. Baker SK (1999) Rumen methanogens, and inhibition of methanogenesis. *Aus J Agri Res* 50: 1293-8.
66. Boadi D, Benchaar C, Chiquette J, Masse D (2004) Mitigation strategies to reduce enteric methane emissions from dairy cows: update review. *Canadian J Anim Sci* 84: 319-35.
67. Ferry JG (1993) *Fermentation of Acetate In: Methanogenesis*, Chapman and Hall: London, UK.
68. Thauer RK (1998) Biochemistry of methanogenesis: a tribute to Marjory Stephenson Prize lecture. *Microbiology* 144: 2377-406.
69. Kong Y, Xia Y, Seviour R, Forster R and McAllister TA (2013) Biodiversity and composition of methanogenic populations in the rumen of cows fed alfalfa hay or triticale straw. *FEMS Microbiol Ecol* 84: 302-15.
70. Marvin-Sikkema FD, Lahpor GA, Kraak MN, Gottschal JC, Prins RA (1992) Characterization of anaerobic fungus from ilama feces. *Microbiol* 138: 2235-41.
71. Jin W, Cheng YF, Mao SY, Zhu WY (2014) Discovery of a novel rumen methanogen in the anaerobic fungal culture and its distribution in the rumen as revealed by real-time PCR. *BMC Microbiol* 14: 104.
72. Orpin CG (1988) Nutrition and biochemistry of anaerobic Chytridiomycetes. *Biosystems* 21: 365-70.
73. Newbold CJ, Lassalas B, Jouany JP (1995) The importance of methanogens associated with ciliate protozoa in ruminal methane production in vitro. *Letters in Applied Microbiology* 21: 230-4.
74. Hackstein JHP (2010) (Endo)symbiotic Methanogenic Archaea (*Microbiology Monographs Book 19*) Springer-Verlag Berlin Heidelberg, Germany.
75. Stams AJ, Plugge CM (2009) Electron transfer in syntrophic communities of anaerobic bacteria and archaea. *Nat Rev Microbiol* 7: 568-77.
76. Minato H, Otsuka M, Shirasaka S, Itabashi H, Mitsumori M (1992) Colonization of microorganisms in the rumen of young calves. *J Gen Appl Microbiol* 38: 447-56.
77. Morgavi DP, Forano E, Martin C, Newbold CJ (2010) Microbial ecosystem and methanogenesis in ruminants. *Animal* 4: 1024-36.
78. Wolin MJ, Miller TL, Stewart CS (1997) Microbe-microbe interactions. In 'The rumen microbial ecosystem'. (Eds PN Hobson, CS Stewart) pp. 467-91.
79. Latham M, Wolin MJ (1977) Fermentation of cellulose by ruminococcus flavefaciens in the presence and absence of Methanobacterium ruminantium. *Appl Environ Microbiol* 34: 297-301.
80. Scheifinger C, Linehan B, Wolin MJ (1975) H₂ production by Selenomonas ruminantium in the absence and presence of methanogenic bacteria. *Appl Microbiol* 29: 480-3.
81. Bryant MP, Small N (1956) Characteristics of two new genera of anaerobic curved rods isolated from the rumen of cattle. *J Bacteriol* 72: 22-6.
82. Dehority BA (1969) Pectin-fermenting bacteria isolated from the bovine rumen. *J Bacteriol* 99: 189-96.
83. Evans PN, Hinds LA, Sly LI, McSweeney CS, Morrison M, et al. (2009) Community composition and density of methanogens in the foregut of the Tammar wallaby (*Macropus eugenii*). *Appl Environ Microbiol* 75: 2598-602.
84. Henderson G, Cox F, Ganesh S, Jonker A, Young W, et al. (2015) Rumen microbial community composition varies with diet and host, but a core microbiome is found across a wide geographical range. *Scientific Reports* 5: 14567.
85. Hegarty RS (1999) Reducing rumen methane emissions through elimination of rumen protozoa. *Aus J Agri Res* 50: 1321-7.
86. Huang XD, Tan HY, Long R, Liang JB, Wright ADG (2012) Comparison of methanogen diversity of yak (*Bos grunniens*) and cattle (*Bos taurus*) from the Qinghai-Tibetan plateau, China. *BMC Microbiology* 12: 237.
87. Jin W, Cheng YF, Mao SY, Zhu WY (2011) Isolation of natural cultures of anaerobic fungi and indigenously associated methanogens from herbivores and their bioconversion of lignocellulosic materials to methane. *Bioresour Technol* 102: 7925-31.
88. Klieve AV, Ouwerkerk D, Maguire AJ, McMillen L (2009) Unusual archaea detected in the foregut of kangaroos. In 'FEMS 2009: 3rd Congress of European Microbiologists'. Pp. 1181. (Federation of European Microbiological Societies).
89. Leahy SC, Kelly WJ, Altermann E, Ronimus RS, Yeoman CJ, et al. (2010) The genome sequence of the rumen methanogen *Methanobrevibacter ruminantium* reveals new possibilities for controlling ruminant methane emissions. *PLoS One* 5: e8926.
90. Moss AR, Jouany JP, Newbold J (2000) Methane production by ruminants: Its contribution to global warming. *Ann Zootech* 49: 231-53.
91. Shin EC, Choi BR, Lim WJ, Hong SY, An CL, et al. (2004) Phylogenetic analysis of archaea in three fraction of cow rumen based on the 16S rDNA sequence. *Anaerobe* 10: 313-9.
92. Skillman LC, Evans PN, Strompl C, Joblin KN (2006) 16S rDNA directed PCR primers and detection of methanogens in the bovine rumen. *Lett Appl Microbiol* 42: 222-8.
93. Skillman LC, Evans PN, Naylor GE, Morvan B, Jarvis GN, et al. (2004). 16S ribosomal DNA- directed PCR primers for ruminal methanogens and identification of methanogens colonising young lambs. *Anaerobe* 10: 277-85.
94. Tajima K, Aria S, Ogata K, Nagamine T, Matsui H, et al. (2000) Rumen bacterial community transition during adaptation to high-grain diet. *Anaerobe* 6: 273-84.
95. Tatsuoka N, Mohammed N, Mitsumori M, Hara K, Kurihara M, et al. (2004) Phylogenetic analysis of methyl coenzyme-M reductase detected from the bovine rumen. *Lett Appl Microbiol* 39: 257-60.
96. Wright ADG, Toovey AF, Pimm CL (2006) Molecular identification of methanogenic archaea from sheep in Queensland, Australia reveal more uncultured novel archaea. *Anaerobe* 12: 134-9.

Submit your next manuscript to Annex Publishers and benefit from:

- ▶ Easy online submission process
- ▶ Rapid peer review process
- ▶ Online article availability soon after acceptance for Publication
- ▶ Open access: articles available free online
- ▶ More accessibility of the articles to the readers/researchers within the field
- ▶ Better discount on subsequent article submission

Submit your manuscript at
<http://www.annexpublishers.com/paper-submission.php>
