ISSN: 2454-499X



Case Report **Open Access**

Tumour or Fungus? A Diagnostic Conundrum

Rajkumar JS1, Nabeel Nazeer*2, Shreya R3, Akbar S2, A Jayakrishna Reddy2 and Anirudh JR2

¹Chairman and Chief Surgeon, LIMA Lifeline Institute of minimal access, Kilpauk, Chennai-600 010 Tamil Nadu, India

²General and Laparoscopic Surgeon, LIMA Lifeline Institute of minimal access, Kilpauk, Chennai-600 010, Tamil Nadu, India

³Final Year General Surgery Resident, Sree Balaji medical college and Hospital, Chrompet, Chennai- 6000 044, Tamil Nadu, India

*Corresponding author: Nazeer N, General and Laparoscopic Surgeon, LIMA Lifeline Institute of minimal access, Kilpauk, Chennai - 600 010, Tamil Nadu, India, Tel: 9176570030, E-mail: nabeelnaslam@gmail.com

Citation: Rajkumar JS, Nabeel N, Shreya R, Akbar S, A Jayakrishna Reddy, et al. (2025) Tumour or Fungus? A Diagnostic Conundrum. J Aids Hiv Infec 10(1): 102

Received Date: March 02, 2025 Accepted Date: March 15, 2025 Published Date: March 19, 2025

Abstract

Candida, a commensal of the oral and the vaginal cavity, sometimes turns invasive, in immunosuppressed individuals. The resultant lesion, known as invasive or aggressive candidiasis, can be extremely refractory to treatment.

This is a case report of an ulcero proliferative lesion in the hard palate that mimicked an intra-oral neoplasm. The clinical picture was confused by the fact that the patient was a chain smoker, and also because the initial MRI images were strongly suggestive of an invasive malignancy. When clinical and radiological signs exactly mimic a malignancy, the picture is confusing.

This is the case report of a 41year old with HIV who was presented with a tumour like ulceroproliferative lesion of the soft palate, with erosion and destruction of the maxilla, and with progression to the base of the skull, treated successfully with long term anti-fungal cocktail.

Keywords: HIV

Introduction

Candida species are human fungal pathogens that cause mucosal and deep tissue infections. There is evidence to suggest that the majority of infections produced by this pathogen are associated with biofilm growth [1]. Immuno compromised patients are prone to opportunistic infections. Lower CD4 count between 200-250 cells/mm3, age and gender are considered as predisposing factors for the increasing incidence rate [2]. The fungal group of infections are caused by Candida albicans (Horn et al., 2009) [1]. Although oropharyngeal candidiasis is a common issue, ranking as a first line sign of HIV to AIDS conversion, only rarely is it invasive, mimicking a tumour in many ways. Invasive candidiasis has an increasing incidence and a crude mortality exceeding

This is a case report of an ulcero proliferative lesion in the hard palate that mimicked an intra-oral neoplasm. The clinical picture was confused by the fact that the patient was a chain smoker, and also because the initial MRI images were strongly suggestive of an invasive malignancy. Fortunately, after multiple biopsies, the hyphae of the candida were identified, and appropriate anti-fungal therapy instituted after which there was a marked diminision in size in the intra oral portion. This case is being presented because of its rarity, and in order to highlight the multiple aspects of aggressive candidiasis in the immuno suppressed.

Case Details

A 41 year old, diagnosed with HIV three years prior to presentation, and on first line Highly Active Antiretroviral Therapy (HAART), presented with a non-healing ulcer of the soft palate. In the last three months, he had lost significant weight, and had a fall of CD4 count by 15%. He also had an increase in viral count, the latter two attributable to noncompliance with therapy for four months.

The ulceroproliferated area on the soft palate was 4 cms across, extending up to the uvula. The edge of the ulcer was everted, and unhealthy slough was seen in the base of the ulcer. The entire lesion was indurated, and bled to examination.

There was a diffuse fullness, pain and tenderness over the left maxilla, and the patient also complained of epistaxis. X-rays showed diffuse thickening of the wall of the maxillary sinus, and suggested involvement of the floor of the antrum. MRI of the skull and sinuses showed infiltration of the posterio inferior part of the left maxilla by a mass lesion. The posterior aspect extended to the base of the skull. Biopsies of the lesion were taken in four quadrants.

There was no histopathological evidence of malignancy, but dense inflammatory infiltrate with macrophages and leukocytes, was found in all four quadrants. Fungal stains revealed no hyphae (Figure 1).



Figure 1: Before

As malignancy was ruled out, a provisional diagnosis of invasive candidiasis was made, and the patient was commenced on Itraconazole, 150 mg/day. The HAART was recommended, ensuring patient compliance this time.

There was a partial improvement of the clinical picture, but not total regression. MRI was repeated 4 months after the administration of Itraconazole, and this showed extension of base of skull disease, but reduction of maxillary disease (Figure 2).



Figure 2: After

An ID consult was sought, and voriconazole was started. In the meantime, he developed palatial anesthesia on the left side, and a suspicion of malignant infiltration of the pterygo palatine fossa was brought up. Accordingly, he underwent multiple biopsies again, which showed the same histopathology as the first time, with no evidence of malignancy or obvious fungal hyphae.

High dose oral Itraconazole (250 mg) and ten day bursts of voriconazole were given for the next three months, and the palatal lesion regressed significantly. A third set of biopsies showed nil malignancy but this time fungal hyphae were obtained from scrapings of the maxillary wall. A fortnight's course of Caspofungin was given, and the patient was then continued on oral Itraconazole.



Figure 3:

At last review, nine months after initial presentation, he still has a residual lesion in the maxilla, but the palatal part has almost completely regressed. His HIV status has stabilized, with a viral count that is negligible and a CD4 count of 690cells/mm³ (Figure 3).

Discussion

Opportunistic infections enter the body due to compromised host defense or through an invasive gateway. Candida infection is more often encountered fungal infection [4]. One third of the HIV positive patients suffer from oral candidiasis [5].

Candida infection develops due to the colonization of the mucocutaneous surfaces. Along with the increased colonization rate HIV is associated with the development of overt disease. During the course of HIV infection, the rate of Candida infection is inversely related to the CD4 counts of the patient which in turn determines the Anti-retroviral treatment [6]. Alterations in the number and differentiation of lymphocytes and dendritic cells play a role in the pathogenesis and evolution of the disease [7].

Candidiasis in HIV disease is no exception, being almost the norm. But invasive/aggressive candidiasis is a different entity, infiltrating surrounding structures, and behaving exactly like a locally invasive malignancy.

This case is educative from many aspects:

The discontinuation of the HAART by the patient probably set off a chain of events, from lowered CD 4 and increased viral counts, triggering off an invasive response by the Candida.

The MRI films were ambivalent, and the invasion of the maxilla and the skull base were highly suggestive of malignancy, warranting multiple biopsies, that were negative for neoplasm. The fact that only the third set of biopsies showed fungal hyphae, points to the innate difficulties of fungus diagnosis. Scrapings from the expanding side of the lesion, will demonstrate fungi, and this was the case here.

This case study is significant for the following reasons – 1. Fungus, especially invasive candiasis can mimic tumour in the immunosuppressed individual. In an era of increasing HIV disease this is an important message. 2. If repeated biopsies of a tumour like lesion turn out to be negative one has to consider rarer conditions like an invasive fungus .3. Avoidance of aggressive procedures in the immunosuppressed groups of patients without a definite tissue diagnosis is a caveat.

Conclusion

Invasive/aggressive Candidiasis, in immune compromised patients, can mimic malignancy. This case report of an Indian patient with aggressive Candidiasis of the palate, that reached the base of the skull, is reported for its unique diagnostic and therapeutic difficulties.

Refrences

- 1. Sardi JC, Scorzoni L, Bernardi T, Fusco-Almeida AM, Mendes MJS (2013) Candida species: current epidemiology, pathogenicity, biofilm formation, natural antifungal products and new therapeutic options. J Med Microbiol 62: 10-24.
- 2. Kumar SK, Gowda SV, Basavarajaiah DM (2013) Incidence of Oral candidiasis among HIV infected patients-Cohort prospective study. Int J Sci Res Pub 3.
- 3. Andes DR, Safdar N, Baddley JW, Playford G, et al. (2012) Impact of Treatment Strategy on Outcomes in Patients with Candidemia and Other Forms of Invasive Candidiasis: A Patient-Level Quantitative Review of Randomized Trials. Clin Infect Dis 54: 1110-22.
- 4. Urs AB, Singh H, Mohanty S, Sharma P (2016) Fungal osteomyelitis of maxillofacial bones: Rare presentation. J Oral Maxillofac Pathol 20: 546.
- 5. McCarthy GM (2012) Host factors associated with HIV-related oral candidiasis. Oral Surgery Oral Med Oral Pathol 73: 181-6.
- 6. Anwar KP, Malik A, Subhan KH (2012) Profile of candidiasis in HIV infected patients. Iran J Microbiol 4: 204-9.
- 7. Romagnoli P, Pimpinelli N, Mori M, Reichart P, Eversole L, Ficarra G (1997) Immunocompetent cells in oral candidiasis of HIV-infected patients: an immunohistochemical and electron microscopical study. Oral Dis 3: 99-105.

Submit your next manuscript to Annex Publishers and benefit from:

- **Easy online submission process**
- > Rapid peer review process
- ➤ Online article availability soon after acceptance for Publication
- ➤ Open access: articles available free online
- More accessibility of the articles to the readers/researchers within the field
- > Better discount on subsequent article submission

Submit your manuscript at http://www.annexpublishers.com/paper-submission.php