

A Simple Method for the Quantification of Atenolol from the Forensic Human Blood and Visceral Samples by High Performance Thin Layer Chromatography

Gayathridevi S*, Sujatha KM, Kruthika N and Krishnaraju SK

Regional Forensic Science Laboratory, Mysuru-570 001, Karnataka, India

*Corresponding author: Gayathridevi S, Regional Forensic Science Laboratory, Mysuru-570 001, Karnataka, India, Tel: +919027385112, Email : math.gayatri@rediffmail.com

Citation: Gayathridevi S, Sujatha KM, Kruthika N, Krishnaraju SK (2023) A Simple Method for the Quantification of Atenolol from the Forensic Human Blood and Visceral Samples by High Performance Thin Layer Chromatography. J Forensic Sci Criminol 11(1): 103

Received Date: September 11, 2023

Accepted Date: September 26, 2023

Published Date: September 30, 2023

Abstract

A simple High-Performance Thin layer chromatography (HPTLC) method has been developed for determination of atenolol in human blood and viscera. The mobile phase was ethyl acetate, acetone, ethyl alcohol, ammonia solution in the ratio of 45:45:7:3. The densitometry scanning at 254 nm was found to be maximum absorption for standard atenolol. The standard and extracted atenolol from samples were detected by HPTLC and confirmed with authenticated spectra at 254 nm. The quantity of the atenolol determined to be 1.6 µg for viscera and 1.5 µg for blood sample.

Keywords: Atenolol; Blood; HPTLC; Viscera

Introduction

Atenolol, 4-(2-hydroxy-3-isopropyl-aminopropoxy) phenylacetamide, is a cardioselective β -blocker used for the treatment of cardiovascular disorders, such as hypertension, angina pectoris, cardiac arrhythmias and myocardial infarction [1]. Hypertension is considered as the crucial health issues in majority of the human population leading to heart attack [2]. Increasing blood pressure causes deaths of about nine million people every year, being responsible for 45% of the deaths due to heart diseases and 51% of strokes [3]. Due to the increased population suffering from hypertension every year, the intake of the β -blockers is increased. Since from three decades the β -blockers are used for treating hypertension among which, Atenolol is considered as most prescribed drug for hypertension [2]. In a single treatment of atenolol drug, the systolic and diastolic blood pressure reduces by 15-20% and also lowers the cardiovascular mortality [4]. As a consequence of the increased utilization of these β -blockers and also different case reports suspected as the deaths caused by the β -blockers were published [5-7]. Hence, several methods have been reported for determining atenolol in human plasma and other biological fluids, including high-performance liquid chromatography (HPLC) [8-14], liquid chromatography - tandem mass spectrometry (LC-MS-MS) [15] capillary electrophoresis (CE) [16] and gas chromatography - mass spectrometry (GC-MS) [17-18]. The benefits of these methods include highly sensitive and selective, and some drawbacks could be: expensive equipment, toxic and expensive solvents (mainly HPLC methods) and usually tedious sample pre-treatment when used for analysing biological samples. Therefore, we report a sensitive and specific HPTLC procedure for determining atenolol from the forensic human blood and viscera samples. The advantages of the present method include simple and single step extraction procedure utilizing inexpensive chemicals and short run time. A simple qualitative and quantitative method was required to analyse and detection of atenolol following oral administration. Current procedure showed high variability due to matrix interference. Hardly were any reports on the determination of atenolol by HPTLC method from the forensic samples. In cases like limited blood draw is required or it is provided with the combination of other drugs which cannot be analysed in the whole blood sample, the atenolol analysis in the whole blood could be necessary. Hence, in this paper we report a simple HPTLC method is reported for the determination of atenolol from the forensic samples of human blood and viscera.

Materials and Methods

Chemicals and Reagents

Atenolol was of commercial grade procured from local Medical stores, manufactured by IPCA laboratories Ltd (50 mg tablet). All the other reagents were of analytical grade and were obtained from Sigma Aldrich. HPTLC aluminium plates, percolates silica gel-60, t_{252} (20X20 cm), 0.2 mm thickness purchased from Merck, Germany.

Blood and Viscera Samples

The blood sample and visceral samples were obtained from corpses auto-spaced in the legal Medical Institute (Mandya district hospital) preserved in saturated solution of NaCl at 4 °C until further use.

Preparation of Standard Solution

Stock standard solution of atenolol was prepared in ethanol at 1mg/mL and maintained at -2°C until and working solution were prepared by diluting with ethanol.

Sample Preparation

Tissue sample: 2 g of tissue was sliced into small fragments and transferred to 250 ml Erlenmeyer flask, to which 100 mL of chloroform was added, shaken for 5-6 h. The organic layer was separated; the aqueous layer was again extracted with 50 mL chloroform. The chloroform layer was separated; all the two organic layers were pooled and passed through anhydrous sodium sulphate over a funnel, evaporated to dryness and used for HPTLC (for instrumentation purpose organic layer should be passed through Florosil column).

Blood sample: 50 mL of post-mortem blood sample was taken in a 250 ml Erlenmeyer flask to which 50 mL of acetonitrile was added, vortexed for 10 min and incubated for 1 h at room temperature. The two solvent layers were mixed and passed through anhydrous sodium sulphate.

Chromatographic Conditions

The samples were applied manually in the form of spots on aluminium plates percolated with silica gel-60 F254 (20X10 cm). Space between spots was 20 mm, slit dimension was 0.45 µm and scanning speed was 20 mm/sec. The mobile phase was ethyl acetate, acetone, ethyl alcohol, ammonia solution in the ratio of 45:45:7:3. The linear ascending chromatogram was developed in a glass chamber saturated with mobile phase, till the mobile phase migrate 8 cm. Following the development, the plates were air dried, spots were visualized under UV lamp-254 nm and densitometric scanning was carried by the scanner (200-400 nm range) 3S-M130319 with Wincat software (CAMAG, Switzerland).

Results and Discussion

The rate of absorption was found to be 50-60% when its orally administered, and the rest is excreted unchanged in the faces. Atenolol undergoes little or no metabolism and is eliminated primarily by renal excretion [8]. Hence atenolol was found to be stable in blood and viscera samples. Spectra for different concentrations of atenolol (2 to 14 µg/ml) were obtained by scanning at 200 to 400 nm wavelength and optimized the symmetrical peaks of the atenolol for accurate and reproducible results. Standard atenolol peak was compared with the atenolol extracted from the post-mortem viscera (Figure 1) and confirmed the peak and wave length at 254 nm. Atenolol extracted from the blood was compared with the standard atenolol spectra (Figure 2). The HPTLC plate (Figure 3) was visualized under the UV light, the sample extracted from blood (Lane b) and viscera (Lane c) were comparable with the standard atenolol (Lane a). Further these spots were confirmed by using spraying reagent Dragandorf (Figure 4). HPTLC coupled with densitometry was well established and could be the best choice for the identification and analysis of drugs in mixtures and in formulations or from biological samples, due to its high-resolution power. It has been successfully applied for the stability studies and the analysis of a vast number of pharmaceuticals [19]. Different wavelengths have been tested (210, 227 and 254 nm), and found to be at 254 nm and gave good scanning results at which peaks were sharp and symmetrical. Minimum noise and considerable sensitivity of the studied compound was obtained. The TLC coupled with densitometric methods could be applied for the determination of atenolol and in different pharmaceutical formulations with acceptable percentage recoveries [20]. HPTLC technique has been explored for the analysis of various drugs [21-22] and pesticides [23-24]. One of the important applications of HPTLC, includes, several samples could be analysed simultaneously, small quantity of mobile phase, reduces the cost and time of analysis and no restrictions on the choice of solvents and mobile phases [21]. Therefore, these methods do not require any extraction procedure always, and could be developed for drug analysis without any interference from excipients [25].

On comparison with HPLC, TLC does not require pure and high concentrated samples. Highly purified sample are required for HPLC columns as their columns affect separation property. The reported method for atenolol analysis in human whole blood and viscera sample is specific and sensitive. Furthermore, this method may be faster for the analysis as it requires very simple method for the preparation of sample, because of which several number of the analysis can be carried in a single day.




Figure 1: Densitometry scanning spectra of Atenolol; **a:** Extracted from Viscera; **b:** Standard Atenolol




Figure 2: Atenolol extracted from blood sample; **a:** Compared with the; **b:** Standard Library




Figure 3: HPTLC plate scanned in UV light at 254 nm; **a:** Standard Atenolol (2 µg) from Commercially Available Sample; **b:** Blood Extract of Deceased; **c:** Visceral Extract of Deceased



Figure 4: Visibility of orange coloured spot when sprayed with Dragandroff reagent; **a:** Blood Sample; **b:** Visceral Sample; **c:** Standard Atenolol

Conclusion

This work provides accurate, sensitive, economic, rapid and selective analytical method for qualitative and quantitative determination of Atenolol from the blood and viscera samples. TLC-Densitometric method for the analysis of atenolol has the advantages over other methods like HPLC, RPHPLC moreover, it can be easily applied in quality control laboratories where economy and time is essential.

Acknowledgment

Director, Directorate of Forensic Science Laboratories, Bangalore and Regional Forensic Science Laboratory, Mysuru are acknowledged for their support.

References

1. Al-Ghannam SMA (2006) A Simple spectrophotometric method for the determination of β -blockers in dosage forms. *J Pharm Biomed Anal* 40: 151-6.
2. Fernandez-Lopez L, Pellegrini M, Rotolo MC, Luna A, Falcon M, et al. (2019) Development and Validation of a Method for the Analysis of Bisoprolol and Atenolol in Human Bone. *Molecules* 24: 2400.
3. World Health Organization (WHO) (2004) The Global Burden of Disease: 2004 Update. World Health Organization, Geneva, Switzerland.
4. Li W, Li Y, Francisco DT, Naidong W (2005) Hydrophilic interaction liquid chromatographic tandem mass spectrometric determination of atenolol in human plasma. *Biomed Chromatogr* 19: 385-93.
5. Mozayani A, Singer P, Jones G (1995) Distribution of metoprolol enantiomers in a fatal overdose. *J Anal Toxicol* 19: 519-21.
6. Fucci N, Offidani C (2000) An unusual death by propranolol ingestion. *Am J Forensic Med Pathol* 21: 56-8.
7. Buajordet I, Ebbesen J, Erikssen J, Brørs O, Hilberg T (2001) Fatal adverse drug events: The paradox of drug treatment. *J Intern Med* 250: 327-41.
8. Spanakis M, Niopas I (2013) Determination of atenolol in human plasma by HPLC with fluorescence detection: validation and application in a pharmacokinetic study. *J Chrom Sci* 51: 128-32.
9. Chatterjee DJ, Li WY, Hurst AK, Koda RT (1995) High-performance liquid chromatographic method for determination of atenolol from human plasma and urine: simultaneous fluorescence and ultraviolet detection. *J Liq Chromatogr* 8: 791-806.
10. Giachetti C, Tencoti A, Canali S, Zanol G (1997) Simultaneous determination of atenolol and chlorthalidone in plasma by high-performance liquid chromatography Application to pharmacokinetic studies in man. *J Chromatogr B* 698: 187-94.
11. Iha MH, Martinez AS, Bonato PS (2002) Enantioselective analysis of atenolol in biologic fluids: comparison of liquid-liquid and solid-phase extraction methods. *J Chromatogr B* 767: 1-9.
12. Miller RBA (1991) Validated high-performance liquid chromatographic method for the determination of atenolol in whole blood. *J Pharm Biomed Anal* 9: 849-53.
13. Morris RG, Saccoia NC, Sallustio BC, Zacest R (1991) Improved high-performance liquid chromatography assay for atenolol in plasma and urine using fluorescence detection. *Ther Drug Monit* 13: 345-49.
14. Chiu FC, Zhang JN, Li RC, Raymond K (1997) Efficient assay for the determination of atenolol in human plasma and urine by high-performance liquid chromatography with fluorescence detection. *J Chromatogr B* 691: 473-7.
15. Li W, Li Y, Francisco DT, Naidong W (2005) Hydrophilic interaction liquid chromatographic tandem mass spectrometric determination of atenolol in human plasma. *Biomed. Chromatogr* 19: 385-93.
16. Arias R, Jimenez RM, Alonso RM, Telez M, Arrieta I, et al. (2001) Determination of the β -blocker atenolol in plasma by capillary zone electrophoresis. *J Chromatogr A* 916: 297-304.

17. Yilmaz B, Arslan S (2009) GC-MS determination of atenolol plasma concentration after derivatization with N-methyl-N-(trimethylsilyl)trifluoroacetamide. *Chromatographia* 70: 1399-404.
18. Amendola L, Molaioni F, Botre F (2000) Detection of beta-blockers in human urine by GC-MS-MS-EI: perspectives for the antidoping control. *J Pharm Biomed Anal* 23: 211-21.
19. Pool CE, Poole SKJ (1989) Progress in densitometry for quantitation in planar chromatography. *Chromatogr B: Biomed Sci Appl* 492: 539.
20. Abdelwahab NS (2010) Determination of atenolol, chlorthalidone and their degradation products by TLC-densitometric and chemometric methods with application of model updating. *Anal Methods* 94: 1994-2001.
21. Parmar VK, Parikh RH, Patel RJ (2016) HPTLC Method for estimation of Topiramate in solubility studies, diffusion studies, plasma, brain homogenate and pharmaceutical formulation. *J Chromatographic Sci* 54: 1105-14.
22. Parmar P, Mehta A (2009) Development and validation of HPTLC method for the estimation of Clotrimazole in bulk drug and tablet formulation. *Indian J Pharm Sci* 71: 451-4.
23. Sanganalmath PU, Nagaraju PM, Sreeramulu K (2019) Determination of quinalphos in human whole blood samples by high-performance thin-layer chromatography for forensic applications. *J Chromatography A* 1594: 181-9.
24. Sanganalmath PU, Nagaraju PM, Sreeramulu K (2020) Development and validation of a simple and rapid thin-layer chromatography-UV densitometry method for the determination of triazophos in human whole blood for forensic toxicological applications. *J Planar Chromatogr-Modern TLC* 2020: 10.1007/s00764-020-00022-3.
25. Patel RB, Patel MR, Bhatt KK, Patel BG (2010) HPTLC method development and validation: Quantification of paliperidone in formulations and in vitro release study. *Analyt Method* 2: 525-31.

Submit your next manuscript to Annex Publishers and benefit from:

- Easy online submission process
- Rapid peer review process
- Online article availability soon after acceptance for Publication
- Open access: articles available free online
- More accessibility of the articles to the readers/researchers within the field
- Better discount on subsequent article submission

Submit your manuscript at

<http://www.annexpublishers.com/paper-submission.php>