

# Point Prevalence of Gastrointestinal Parasites in Animals at a Zoological Garden in Ibadan, Oyo State, Nigeria

Ojo PO\*, Adediran OA and Osundara OO

Department of Veterinary Parasitology and Entomology, Faculty of Veterinary Medicine, University of Ibadan, Oyo State, Nigeria.

\*Corresponding Author: Ojo PO, Department of Veterinary Parasitology and Entomology, Faculty of Veterinary Medicine, University of Ibadan, Oyo State, Nigeria, E-mail: philipsorji@gmail.com

**Citation:** Ojo PO, Adediran OA, Osundara OO (2026) Point Prevalence of Gastrointestinal Parasites in Animals at a Zoological Garden in Ibadan, Oyo State, Nigeria. J Vet Sci Ani Husb 14(1): 104

**Received Date:** June 12, 2026 **Accepted Date:** June 23, 2026 **Published Date:** June 29, 2026

## Abstract

Gastrointestinal parasites are a serious impediment to animal production and sustainability in Nigeria. This study investigates the prevalence of gastrointestinal parasites among captive animals in the Zoological Garden in Ibadan, Nigeria. A total of 57 freshly voided feces were collected from seven species of zoo animals. The overall prevalence of helminth and protozoal parasites was 27 (47.5%) and 28 (49.1%), respectively. Carnivorous species had the highest prevalence of *Toxocara* species (100%). Hyenas had a high prevalence of multiple parasites, including *Capillaria* species (100%), *Toxocara* species (66.7%), *Taenia* species (66.7%) and *Cryptosporidium* species (33.3%). Primates exhibited high protozoan infection rates (44.4%) and lower helminth infection rates (27.8%). However, hoofed mammals showed higher helminth prevalence (63.7%) than protozoan parasites. Equine species had multiple helminth infections with the highest prevalence in *Anoplocephala* species (80%), followed by *Fasciola* species (60%), *Strongyloides* species (60%) and *Dicrocoelium* species (40%). In this study, there was co-infection with multiple helminths in camel. Coccidian oocysts and helminth burdens in avian species were (53.3%) and (13.3%), with *Eimeria* species, *Cryptosporidium* species and *Heterakis gallinarum* having a prevalence of 46.6%, 40% and 13.3%, respectively. This research demonstrated high prevalence of gastrointestinal and zoonotic parasites, including *Cryptosporidium* species, *Strongyloides* species and *Toxocara* species across different host groups, highlighting the potential risk to public health.

**Keywords:** Prevalence; gastrointestinal parasites; captive; wild animals

## Introduction

Nigeria is home to several natural beauties and has a rich, diverse wildlife. Several factors threaten the survival of wild animals, including habitat loss, poaching, climate change, and wildlife diseases, especially those caused by gastrointestinal parasites [1]. Zoological gardens display collections of wild animals for research, conservation, education, and recreation. Animals in zoos are captured from their natural habitats and kept in man-made environments [2, 3]. The main challenges of keeping wild animals in captivity are parasitic diseases [4]. In their natural habitats, wild animals may have immunity to ectoparasites and endoparasites, or maintain a balance with them as they roam across large areas. Parasites can infect hosts, but the animal's immune system determines whether disease develops [5]. Capturing animals for captivity can affect their overall welfare and ecological needs, increasing their vulnerability to diseases [6], even when they are provided with adequate food, water, and hygiene in zoological gardens [7]. When pathogens spread between species, they can disrupt ecosystems by reducing biodiversity, changing the structure and behavior of animal populations, and affecting interspecies interactions, including predator-prey relationships [8]. Gastrointestinal parasites, which inhabit the digestive tract, are commonly detected through faecal analysis. Because their presence, absence, or relative abundance reflects the condition of host species, these parasites serve as powerful bio indicators of ecosystem health [9]. Monitoring parasites provides insight into the stability of animal populations, the resilience of ecological communities, and the potential impacts of environmental changes. In this way, parasites are not merely agents of disease but also valuable markers that help assess the balance and integrity of ecosystems. Therefore, this study aimed to determine the prevalence of gastrointestinal parasites in a zoological garden at Ibadan, Nigeria.

## Materials and Methods

### Study site


The Zoological Garden is situated at a latitude of 7°23'47"N and longitude 3°55'0"E in Ibadan, Oyo State, Nigeria. It was founded in 1948 as a menagerie for teaching and research. The climate in Ibadan is tropical, with clear wet and dry seasons, supporting a diverse range of animal life. To prevent parasitic diseases, zoo animals at this facility are treated routinely and whenever tests are positive for parasitic infections. Veterinarians and attendants carry out regular examinations on the animals in the Zoological Garden. When diarrhoea is observed, faecal examination is performed, and any positive animals are treated accordingly, with supplementary analysis performed after the treatment period. Furthermore, in this zoo, newly acquired animals are quarantined and clinically examined, with faecal and blood samples collected and analysed by veterinarians to determine their health status.

### Sample collection and parasitological analysis

Faecal samples were collected in the morning with the assistance of the Zoo animal handlers. Individual freshly voided faecal samples were collected. A total of 57 samples were collected from seven species of wild animals housed in the Zoological Garden. Samples were collected four to six weeks after the animals' routine deworming. Of these samples, Felines accounted for 10, canines (3), equines (5), Camel (1), Primates (18), Ruminants (5) and Aves (15). All samples were stored at 4°C, transported to the laboratory, and examined within 48 hours. Samples were subjected to macroscopic examination to verify the presence of adult nematodes, cestodes and parasite fragments and to microscopic examination using flotation and sedimentation techniques. Fresh wet-mount smears, along with modified Ziehl-Neelsen and Giemsa-stained smears prepared from 10 mL of sediment obtained from faecal samples processed via formol-ether centrifugation, were also prepared. Microscopically, samples were evaluated at magnifications of 100x, 400x, and 1,000x.

## Results

The overall parasite burden among animals in the Zoological Garden was high with helminths (47.5%) and protozoa (49.1%) of the total samples examined. Felines and canines exhibited the highest helminth prevalence (100%). Protozoan infections were also common in these groups, affecting 50% of felines and 33.3% of canines (Figure 1). Primates exhibited lower helminth incidence (27.8%) but relatively high protozoan prevalence (44.4%). Hoofed mammals had a mixed burden, with helminths detected in 63.7% and protozoa in 45.5%. Avian species exhibited the lowest helminth prevalence (13.3%) but the highest protozoan burden (53.3%).



**Figure 1:** Over all Helminth and Protozoan incidences among animals in the University of Ibadan Zoological Garden

Among the carnivores, all lions (100%) were infected with *Toxocara* spp. and Hyaenas were also heavily parasitised by *Capillaria* spp. (100%), while *Taenia* spp. and *Toxocara* spp. were present in 66.7% and *Cryptosporidium* spp. at 33.3%. Equines exhibited a diverse parasite spectrum, with high prevalence of *Anoplocephala* spp. (80%), followed by *Fasciola* spp. (60%), *Cryptosporidium* spp. (60%), *Strongyloides* spp. (60%) and coccidian oocysts at 20%. Multiple parasites were detected in Camel, including *Dicrocoelium* spp., *Dictyocaulus* spp., *Anoplocephala* spp., and *Trichuris* spp. In primates, protozoan infections were slightly higher with Coccidian oocysts (38.9%), *Cryptosporidium* spp. (27.8%) and *Strongyloides* spp. at 27.8%. Ruminants showed high prevalence of Coccidian oocysts (40%) and *Cryptosporidium* spp. (20%), while *Fasciola* spp. had 20% prevalence. Birds exhibited the highest protozoan burden, with Coccidian oocysts (46.6%) and *Cryptosporidium* spp. (40%). Helminth infection was relatively low, with *Heterakis* spp. detected in 13.3% of the aviary (Table 1).

**Table 1:** Prevalence of gastrointestinal parasites among captive animals in the University of Ibadan Zoological Garden

Parasites	Lions (N=10) %	Hyaena (N=3) %	Equine (N=5) %	Camel (N=1) %	Primates (N=18) %	Ruminants (N=5) %	Aves (N=15) %
<i>Toxocara</i> spp	10 (100)	2 (66.7)	-	-	-	-	-
<i>Capillaria</i> spp	-	3 (100)	--	-	-	-	-
<i>Taenia</i>	-	2 (66.7)	-	-	-	-	-
<i>Strongyloides</i> spp	-	-	3 (60)	-	5 (27.8)	--	-
<i>Fasciola</i> spp	-	-	3 (60)	-	-	1 (20)	-
<i>Dicrocoelium</i> spp	-	-	2 (40)	1 (100)	-	-	-
<i>Dictyocaulus</i> spp	-	-	-	1 (100)	-	-	-
<i>Anoplocephala</i> spp	-	-	4 (80)	1 (100)	-	-	-
<i>Heterakis</i> spp	-	-	-	-	-	-	2 (13.3)

Trichuris spp	-	-	-	1 (100)	-	-	-
Cryptosporidium spp	3 (30)	1 (33.3)	3 (60)	-	5 (27.8)	1 (20)	6 (40)
Coccidian oocysts	2 (20)	-	1 (20)	-	7 (38.9)	2 (40)	7 (46.6)

- = Negative

Several zoonotic gastrointestinal parasites were identified across different animal groups in the University of Ibadan Zoological Garden (Table 2). *Cryptosporidium* spp. was the most widely distributed and detected in felines, hoofed mammals, primates, and birds. *Strongyloides* spp. was observed in hoofed mammals and primates and *Toxocara* spp. was restricted to felines and canines. *Fasciola* spp. was detected only in hoofed mammals and *Taenia* spp. was observed in canines.

Animals	<i>Cryptosporidium</i> spp	<i>Strongyloides</i> spp	<i>Toxocara</i> spp	<i>Fasciola</i> spp	<i>Capillaria</i> spp	<i>Taenia</i> spp
Feline	+	-	+	-	-	-
canine	+	-	+	-	+	+
Hoofed mammals	+	+	-	+	-	-
Primates	+	+	-	-	-	-
Aves	+	-	-	-	-	-

= Negative, + = Positive

Among birds, *Heterakis* spp. was detected in speckled pigeons and White stork goose, while *Cryptosporidium* spp. was present in White stork goose and Senegal parrots. Coccidian oocysts were widespread, occurring in speckled pigeons, African grey parrots, White goose, and White stork goose. Overall, protozoan infections were more common than helminths in avian species. However, no parasite was found in Vultures and Marabou storks (Table 3). In non-human primates, *Cryptosporidium* spp. was the most prevalent in Mona monkeys, Green monkeys, and Chimpanzees. *Strongyloides* spp. was observed in Green monkeys, while Coccidian oocysts were found in Mona monkeys, Green monkeys and Chimpanzees.

**Table 3:** Parasites detected in various species of birds and Non-Human Primates at the University of Ibadan Zoological Garden

Animal groups	Parasites			
	<i>Heterakis</i> spp.	<i>Cryptosporidium</i> spp.	<i>Strongyloides</i> spp.	Coccidian oocysts
Vulture	-	-	-	-
Speckled Pigeon	+	-	-	+
African grey Parrot	-	-	-	+
Marabou stork	-	-	-	-
White goose	-	-	-	+
White stork Goose	+	+	-	+
Senegal Parrot	-	+	-	+
Non-Human Primates				
Mona Monkeys	-	+	-	+
Baboons	-	-	-	-

Green Monkeys	-	+	+	+
Chimpanzees	-	+	-	+

- = Negative, + = Positive

In felines, adult lions were positive for *Toxocara* spp. Adult lions also harboured Coccidian oocysts and *Cryptosporidium* spp. Hyenas exhibited a broader parasite spectrum, with *Toxocara* spp., *Cryptosporidium* spp., *Capillaria* spp., and cestodes detected. Horses were infected with *Cryptosporidium* spp., *Anoplocephala* spp., and *Strongyloides* spp., while donkeys carried Coccidian oocysts, *Cryptosporidium* spp., and *Anoplocephala* spp. The camel was positive for mixed helminth infections including *Dictyocaulus* spp. (lungworm), *Dicrocoelium* spp., *Anoplocephala* spp and *Trichuris* spp. (Table 4).

**Table 4:** Parasites detected in Feline, Canine, Equine, Ruminants and Camels at the University of Ibadan Zoological Garden

Animals	<i>Toxocara</i> spp.	Coccidian oocysts	<i>Cryptosporidium</i> spp.	<i>Capillaria</i> spp.	<i>Anoplocephala</i> spp.	<i>Strongyloides</i> spp.	<i>Dictyocaulus</i> spp.	<i>Fasciola</i> spp.	<i>Taenia</i> spp.	<i>Trichuris</i> spp.	<i>Dicrocoelium</i> spp.
Adult Lions	+	+	+	-	-	-	-	-	-	-	-
Hyenas	+	-	+	+	-	-	-	-	+	-	-
Horses	-	-	+	-	+	+	-	-	-	-	+
Donkeys	-	+	+	-	+	-	-	-	-	-	-
Camel	-	-	-	-	+	-	+	-	-	+	+

- = Negative, + = Positive

## Discussion

Gastrointestinal parasites are undoubtedly the bane of livestock and wildlife productivity and sustainability in Nigeria. This study demonstrated a high prevalence of gastrointestinal parasites among captive animals at the University of Ibadan Zoological Garden, with overall burdens of helminths at 47.5% and protozoa at 49.1%. The high prevalence observed in this research highlights captivity as a key risk factor for parasite transmission driven by confinement, environmental contamination, and possible interspecies contact [10]. It also aligns with previous reports that parasitic infections are widespread in zoological settings, particularly in tropical regions where environmental conditions favor parasite survival and transmission [11].

Carnivores exhibited the highest helminth burden (100%), particularly *Toxocara* spp. in felines and canines, reinforcing the well-established host specificity and transmission efficiency of ascarid infections in wild carnivores. Similar patterns have been reported in captive zoo carnivores, where persistent environmental contamination and feeding of raw meat contribute to sustained transmission cycles [10]. The concurrent detection of protozoa in these groups further indicates exposure to contaminated water or fomites, as protozoan cysts are environmentally resistant and are readily transmitted in captive settings. Similar studies in captive lions have reported high *Toxocara* prevalence linked to feeding practices involving raw meat and inadequate parasite control [12]. The presence of protozoa in these groups further indicates exposure to contaminated water or food, as protozoan cysts such as *Cryptosporidium* spp. are known for their environmental persistence [13].

In contrast, primates showed lower helminth prevalence but relatively high protozoan infection rates, supporting earlier observations that protozoan infections dominate in non-human primates under captive conditions [14] and also consistent with recent molecular studies demonstrating the predominance of protozoan parasites such as *Cryptosporidium* spp. and *Giardia duodenalis* in captive primate populations [15]. This pattern likely reflects faecal-oral transmission facilitated by social behavior, enclosure sharing, and hygiene limitations. The detection of *Cryptosporidium* spp. and coccidian oocysts among several primate species is particularly important given their zoonotic potential and documented transmission between primates and hu-

mans [16]. However, the absence of detectable parasites in Baboons may reflect effective immune responses, better enclosure hygiene, or recent antiparasitic treatment.

Hoofed mammals showed a mixed parasite burden, with higher helminth prevalence (63.7%) consistent with grazing-related exposure to helminth infective stages in contaminated pastures or feeds and water. Comparable findings have been reported in zoo ungulates, where pasture contamination and the availability of intermediate hosts sustain helminth life cycles [17]. The detection of trematodes and cestodes further supports the role of environmental and ecological factors, including intermediate hosts, parasite transmission within captive systems.

Avian species exhibited lower helminth prevalence but higher protozoan burdens, indicating that protozoan infections, particularly coccidia and *Cryptosporidium*, dominated in birds under captive conditions. This aligns with a report showing that protozoan parasites are more prevalent in avian species due to high stocking densities, stress-induced immunosuppression, and rapid environmental dissemination of infective oocysts [18]. The widespread occurrence of coccidian oocysts across multiple bird species suggests persistent contamination of enclosures; the absence of parasites in some species may reflect species-specific resistance or lower exposure [10]. The absence of detectable parasites in some bird species may reflect species-specific resistance or differences in exposure and enclosure hygiene.

The widespread occurrence of zoonotic parasites, particularly *Cryptosporidium* spp., underscores the public health implications of these findings. Recent molecular epidemiological study confirms the broad host range and cross-species transmission potential of *Cryptosporidium* in zoo settings, highlighting its importance in One Health frameworks [19]. The detection of other zoonotic helminths, including *Strongyloides* spp. and *Toxocara* spp., further emphasises the risk of environmental contamination and human exposure, particularly for animal handlers and visitors.

Overall, the high prevalence of gastrointestinal parasites observed in this study, within a month and a half after routine deworming, indicates that the animals' enclosure is continually seeded with helminth ova. Improved parasite control strategies are needed in zoological settings. Regular faecal monitoring, strategic deworming, enhanced sanitation, and water quality management are essential to reduce infection pressure. Importantly, the presence of zoonotic parasites reinforces the need for integrated One Health approaches that consider the interconnected health of animals, humans, and the environment.

## Conclusion

The study revealed a high prevalence of gastrointestinal parasites among captive animals. Carnivores showed the highest helminth infections, while birds and primates had more protozoan infections, likely due to differences in diet, hygiene, and environmental exposure. Several gastrointestinal parasites identified include *Toxocara* spp., *Cryptosporidium* spp., *Strongyloides* spp., and *Fasciola* spp., indicating complex transmission dynamics, with zoonotic species posing potential public health risks. Overall, the findings reinforce the need for improved hygiene, routine parasite monitoring, and effective deworming strategies to enhance animal health and reduce public health risk.

## Conflicts of interest

Authors declare no conflicts of interest exist.

## References

1. Jenkins EJ, Simon A, Bachand N, Stephen C (2015) Wildlife parasites in a One Health world. *Trends in Parasitology*. 31: 174–180.
2. Mir AQ, Dua K, Singla LD, Sharma S, Singh MP (2016) Prevalence of parasitic infection in captive wild animals in Bir Moti Bagh mini zoo (Deer Park), Patiala, Punjab. *Vet World*. 2016. 9: 540-3.
3. Schulte-Hostedde AI and Mastromonaco GF (2015) Integrating evolution in the management of captive zoo populations. *Evolutionary Applications*. 8: 413-422.
4. Borghare AT, Bagde VP, Jaulka AD, Katre DD, Jamde PD, et al. (2009) Incidence of gastrointestinal helminthiasis of captive wild deer at Nagpur. *Veterinary World* 2: 337-338.
5. Raja MMRU, Dey AR, Begum N, Kundu UK, Ashad FA (2014) Coprological prevalence of gastrointestinal parasites in carnivores and small mammals at Dhaka Zoo, Bangladesh. *J Threat Taxa* 6: 5574-5579.
6. Goossensa E, Dornya P, Boomkerd J, Vercammen F (2005) A 12-month survey of the gastrointestinal helminths of antelopes, gazelles and giraffes kept at two zoos in Belgium. *Veterinary Parasitology*. 127: 303-312.
7. Dhakal P, Sharma HP, Shah R, Thapa PJ, Pokheral CP (2023) Copromicroscopic study of gastrointestinal parasites in captive mammals at Central Zoo, Lalitpur, Nepal. *Vet Med Sci*. 9: 457-464.
8. Hillary S, Young IM, Parker GS, Gilbert AS, Guerra CL (2017) Introduced Species, Disease Ecology, and Biodiversity–Disease Relationships. *Trends in Ecology and Evolution*. 32: 41-54.
9. Maharjan B, Jain P, Koju NP (2025) Zoonotic risks and conservation challenges: Gastrointestinal parasites in wild mammals of Chitwan National Park, Nepal. *International journal for parasitology. Parasites and Wildlife*. 26: 101041.
10. Cai W, Zhu Y, Wang F, Feng Q, Zhang Z, et al. (2024) Prevalence of Gastrointestinal Parasites in Zoo Animals and Phylogenetic Characterisation of *Toxascaris leonina* (Linstow, 1902) and *Baylisascaris transfuga* (Rudolphi, 1819) in Jiangsu Province, Eastern China Animals.
11. Ferdous S, Chowdhury J, Hasan T, Dutta P, Rahman MM, et al. (2023) Prevalence of gastrointestinal parasitic infections in wild mammals of a safari park and a zoo in Bangladesh. *Vet Med Sci*. 9: 1385-1394.
12. Mukarati NL, Vassilev GD, Tagwireyi WM, Tavengwa M (2013) Occurrence, prevalence and intensity of internal parasite infections of African lions (*Panthera leo*) in enclosures at a recreation park in Zimbabwe. *J Zoo Wildl Med* 44: 686-93.
13. Xiao L (2010) Molecular epidemiology of human cryptosporidiosis: An overview. *Parasitology*, 137: 1181–1196.
14. Nizeyi JB, Mwebe R, Nanteza A, Cranfield MR, Graczyk T (2002) Cryptosporidiosis in people sharing habitats with free-ranging mountain gorillas (*Gorilla gorilla beringei*) in Uganda. *American Journal of Tropical Medicine and Hygiene*, 66: 442–444.
15. Karim MR, Li J, Rume FI, Sumon SMR, Selim ASM, et al. (2021) Occurrence and molecular characterisation of *Cryptosporidium* spp. and *Giardia duodenalis* among captive mammals in the Bangladesh National Zoo. *Parasitology International*. 84: 102414

- 
16. Ryan U, Zahedi A, Paparini A (2016) *Cryptosporidium* in humans and animals a one health approach to prophylaxis. *Wiley Parasite Immunology* 38: 535 – 547.
17. Bhat RA, Tak H, Bhat BA, Dar JA, Ahmad R (2022) Gastrointestinal helminth parasites of wild ungulates in Hirpora Wildlife Sanctuary, Kashmir, India. *J Parasit Dis.* 46: 804-810.
18. Permin A, Hansen JW (1998) Epidemiology, diagnosis and control of poultry parasites. *FAO Animal Health Manual No. 4.* Food and Agriculture Organisation of the United Nations, Rome.
19. Zhao Q, Pei Z, He Y, Jia T, Zhang Y, et al. (2025) Expansion of the known host range of *Giardia duodenalis* and *Cryptosporidium* spp. in captive wildlife at Beijing Zoo. *Parasite.* 32: 4.

Submit your next manuscript to Annex Publishers and benefit from:

- ▶ Easy online submission process
- ▶ Rapid peer review process
- ▶ Online article availability soon after acceptance for Publication
- ▶ Open access: articles available free online
- ▶ More accessibility of the articles to the readers/researchers within the field
- ▶ Better discount on subsequent article submission

Submit your manuscript at

<http://www.annexpublishers.com/paper-submission.php>