

Establishing a Special Open Field Test Appliance for Tree Shrews Evaluates their Stressed Locomotor Behavior

Chuan He^{1,2#}, Jing Dai^{1#}, Minjie Zhou^{1#}, Meng LV³, Jing Wang^{*1}, Jun Gao^{*1,2}

¹Department of Neurobiology, School of Basic Medical Sciences, Nanjing Medical University, Nanjing 211166, Jiangsu, China

²Department of Rehabilitation Medicine, Jiangsu Shengze Hospital Affiliated to Nanjing Medical University, Suzhou, Jiangsu 215228, China

³Animal Core Facility of Nanjing Medical University, Nanjing Medical University, Nanjing 211166, Jiangsu, China

[#]These authors contributed equally to this work.

***Corresponding author:** Jun Gao, Professor, Doctoral Tutor, Department of Neurobiology, School of Basic Medical Sciences, Nanjing Medical University, Nanjing 211166, Jiangsu, China, Tel: 86-025-86869347, E-mail: gaojun@njmu.edu.cn and Jing Wang, Doctor, Lecturer, Department of Neurobiology, School of Basic Medical Sciences, Nanjing Medical University, Nanjing 211166, Jiangsu, China, Tel: 86-025-86869432, E-mail: wangjing@njmu.edu.cn

Citation: Chuan He, Jing Dai, Minjie Zhou, Meng LV, Jing Wang, et al. (2020) Establishing a Special Open Field Test Appliance for Tree Shrews Evaluates their Stressed Locomotor Behavior. J Vet Sci Ani Husb 8(1): 104

Received Date: May 02, 2020 **Accepted Date:** August 11, 2020 **Published Date:** August 13, 2020

Abstract

It has a great significance to establish a high repetitive and special system to evaluate locomotion for tree shrew animal models, especially neuropsychiatric models. Thus, we designed a special open field test appliance for tree shrews by using a polymer PMMA board (plexiglass). The locomotion of naïve tree shrews was detected twice in it. By using Any-maze Software tracing the motion trajectory, we found that the motion trajectory and the distance were very similar in both tests. It demonstrated that the appliance designed for tree shrews had extremely high reliability and stability. Then we used this system to evaluate the activity of the stressed tree shrews, and observed the reduced activity of the tree shrews after a 2-week light-stress. Consistent with this, the ACTH level in the urine of the stressed tree shrews was enhanced. Taken together, the light stress could lead to a significant reduction of locomotion which could be investigated by our open field experiment system. Furthermore, the data from our special open field experiment system is stable and accurate for testing the locomotion behavior of tree shrews.

Keywords: Tree Shrews; Open Field Test; Stress; Motion Trajectory

List of abbreviations: OFT: Open Field Test; ACTH: Adrenocorticotrophic Hormone; ELISA: Enzyme-Linked Immunosorbent Assay; ICC: Intra-group Correlation Coefficient; HPA: Hypothalamic Pituitary-Adrenal; CRH: Corticotropin-releasing Hormone; PVN: Paraventricular Nucleus

Introduction

In recent years, the role of stress-induced neuroendocrine system changes in the pathogenesis has attracted much attention. Stress caused by abominable and demanding conditions can cause different biological responses according to different stress conditions. It is generally believed that transient and moderate stress can trigger adaptive responses, such as a 'fight or flight response', and promote adaptation and resilience to stress. Conversely, chronic or excessive stress can lead to cognitive and affective dysfunction, and can easily lead to mental disorders [1-5]. There are many ways to establish animal stress models depending on the purpose of the study. Among the many methods for creating models, the closest method to human stress is the animal stress modelling method [6]. The tree shrews belong to the sister of primates [7]. Its shape is small, and the whole body is fluffy similar to the squirrel. It was found that the tree shrews had timid and docile characters. They are very sensitive to psychological stress. Stress from bondage can lead to the increased cortisol in tree shrews. The stress depressed tree shrew model showed that there was a continuous increase in cortisol level of subordinate tree shrews, which could lead to atrophy of hippocampal CA3 neurons. This phenomenon is consistent with the atrophy of the hippocampus in patients [8]. Another important index in the evaluation of the stress model inactivity is the open field test. Open field test (OFT), which is also called an open box experiment, is widely used in the study of neuropsychiatric diseases represented by stress behavior. However, most of the current OFT boxes are designed for rodents, because of the large differences between tree shrews and rodents in their living habits, which is not suitable for evaluating the activity of the tree shrews. Moreover, the activity characteristics of the tree

shrews are mainly jumping, while the OFT box of the existing rodents is selected to evaluate their activity, the data of the vertical plane will be not true enough. Recently, most of the locomotor evaluation has been conducted in the home-cages of tree shrew. Although the problem of authenticity can be solved, the background is too complex, which can cause the problem of poor tracking and data loss. Here, we designed a specific open-field laboratory for tree shrews to help us to get the reliable behavior data of tree shrews conveniently. By it, we did the OFT twice, and assessed the accuracy and stability of the data of the tree shrews' behavior experiment. Moreover, to investigate its sensitivity, we investigated the effects of mild stress on the activities of tree shrews by our designed equipment. And we observed the alteration of the activities in the tree shrews induced by light-stress for 2 weeks through this system.

Materials and Methods

Experimental Animals

Since the female tree shrews have menstruation, we only used male tree shrews in the present experiments to avoid the interference of it. Moreover, we wanted to observe the effect of stress on male tree shrews' behavior by self-control (before and after stress) and to investigate the stability of our new OFT instrument. Thus, in the experiments, 15 male adulthood (aged 1-2 years, weighted 120-160 g, and fed with fresh cooked porridge which materials are bought from the Kunming institute of zoology of China) Chinese tree shrews were selected. The tree shrews were provided by the Kunming Institute of Zoology, Chinese Academy of Sciences, and the tree shrews were raised at the Animal Core Facility of Nanjing Medical University. All animals were given ad libitum access to food and water and were housed under a 12-h light/dark cycle (house light on from 8:00 am to 8:00 pm). According to local standards in Yunnan province, the working illumination of tree shrews' rooms is set to 150-300 lux, and that of the area to 50-100 lux. All animal experiments were performed following the recommendations of the Experimental Animal Ethics Committee at Nanjing Medical University.

Methods

Preparation of Special OFT Box for Tree Shrews: The open-field experimental process and equipment for rodents have been accurately designed, but those for tree shrews are lacked. At present, most of researchers use tree shrews' breeding cages or open field experiment boxes for rats to do tree shrews' behavior test. The biological behavior between rats and tree shrews is very different, and the obtained data is inconsistent with the living habits of tree shrews to a certain extent. The repeatability and stability of the data of multiple experiments are low. Therefore, to set up a suitable behavioral experiment box for tree shrews has become the focus of this research.

According to the relevant literature [9,10] and combining with the characteristics of tree shrews, we designed the special OFT box for tree shrews (Figure 1A). The special material is a kind of polymer PMMA polymethyl methacrylate sheet (plexiglass), which is different from the traditional materials for OFT box for the rodents (Table 1). Polymer PMMA polymethyl methacrylate (PMMA) plate has a crystal-like transparency with a light transmittance of more than 92%, which is helpful to capture the image without disturbing the tree shrew experiment. Moreover, PMMA has a light transmittance comparable to glass, but its density is only half of that of glass, which is convenient for frequent handling of box top cover in the process of behavioral operation. Furthermore, it is not as fragile as glass, and even if it is damaged, there will be no sharp fragments like glass, which can reduce the risk of injury during the operation of the experimenter. The size of the box is 50 cm × 50 cm × 80 cm (length × width × height) for tree shrews, which is also different from the OFT box for the rodents (Table 2). Considering the tree shrews' habits from the instinctive behavior in the wild environment, the beam is designed at a height of 40 cm (Table 2) which is made of the frosted material to prevent slippage and make it easier for tree shrews to move autonomously in the direction of the ground. Moreover, the frosted material is easy to be cleaned and has a high degree of safety. The open-field box is composed of a clear plexiglass top, detachable fencers, a recessed white bottom plate and a frosted clear plexiglass rod running through the box. Moreover, in the rodents, we use a central area to evaluate the anxiety-like behavior in the open field test, while the alteration in jumping was used in tree shrews [11]. In the present experiments, the open-field box is divided into three parts from top to bottom: upper part, middle part, and down part (Figure 1B). In the behavior test, both the total distance and the changed activity of the tree shrews in the middle or upper part would be investigated.

Materials	Skid Resistance	Durability	Anti- Alcohol	Transmission of Light	Odour Residue
PMMA	√	√	√	√	√
Iron	×	√	√	×	√
Wood	√	×	×	×	×
Nylon	√	√	√	×	√
Engineering Plastics	√	√	×	√	√

Table 1: Comparison of properties of different materials

	Size (cm)	Hierarchy
For Tree Shrew	50 × 50 × 80	Yes
For Mice	50 × 50 × 30	No
For Rats	100 × 100 × 40	No

Table 2: The difference in the structure of the open field box between rodents and tree shrews




Figure 1: Special OFT box for tree shrews (A) The side view of an open field box; (B) The front view of the open field box and the distribution patten of the OFT box

Open Field Test: Open field test (OFT) is a common method to access exploratory behavior and general activity in rodents [12,13]. It has a lot of use in physiological and pathological conditions, including but not limited in the neuropsychiatric disease represented by stress behavior [14]. A camera was set in front of the transparent plexiglass front baffle on the front of the box to monitor the activity of the tree shrews. Before the experiment, the door of the sleeping box was opened. The tree shrews entered the box through the round hole by themselves, avoiding the stress of grasping on them. The door of the sleeping box was closed quickly and video recording was started. The tree shrews were allowed to move freely for 15 minutes, and the movement parameters of the tree shrews in the horizontal and vertical directions, the average speed and the number of jumps were recorded respectively during the experiment. At the end of the experiment, the door of the sleeping box was opened. After the tree shrews entered the sleeping box through the round hole, the door was closed quickly. After removing the sleeping box, we successively removed the top cover with the air hole, front baffle on the front, frosted glass rod through the box and detachable silicone non-slip pad, and then washed and dried with water. After wiping the residual urine and faeces in the tank with flat paper, spray 70% alcohol to eliminate the residual odour of the experimental tree shrews. The whole operation process eliminates the need to grab the tree shrews and eliminates the artificial stress caused by grasping the tree shrews. Therefore, the most basic activity and emotional state of the tree shrews can be objectively evaluated.

Selection of Detection Software for Tree Shrews' OFT System: The Any-maze Animal Behavior Video Analysis System (ST-60000, Sterling, USA) is a scientific analysis software that provides an advanced grayscale recognition algorithm that adjusts the threshold when the animal's colour contrasts with the background colour. It's easy to lock the target and accurately track the trajectory of the animal's movement with powerful analytical statistics function. We used this system to record the animal's original activity video, the activity track, and get the real-time changes in animal activity patterns and the animal trajectory data.

Stress Experimental Process: We extended the normal circadian light by four hours at night to simulate the pattern of staying up late in humans. Fifteen naïve tree shrews were tested in the open field test (OFT) at the end of the first week and the end of the second week. The Any-maze software was used to track the locomotion. Then, the house light being on was prolonged by 4 hours. That is the house lights on from 8:00 am to 12:00 pm. After 2 weeks of treatment, the tree shrews' locomotion was detected by OFT again. What's more, the concentration of adrenocorticotrophic hormone (ACTH) in the urine of tree shrews before and after stress was detected by enzyme-linked immunosorbent assay.

Enzyme-Linked Immunosorbent Assay: The alteration of ACTH is the index for stress [12]. Thus, the urine samples of tree shrews were collected over dry ice in tubes before breakfast between 07:45 and 08:00. Urine samples were stored at -20 °C until analysis. The concentration of corticotrophin in urine before and after stress was measured by enzyme-linked immunosorbent assay (ELISA). It was measured using an ELISA kit commercially available according to the manufacturer's instructions (ACTH, ml001895, mlbio, China).

Statistical Methods

Statistical analysis was performed by using SPSS 19.0 software, and statistical mapping was performed by using Origin Pro 8.5 software. The data were presented as means \pm SEM. The differences between the two groups were compared via a two-tailed Student's *t*-test. $p < 0.05$ was considered statistically significant.

Results

Tree Shrews' Special OFT Box can Stably Detect Tree Shrews' Activity

After putting the tree shrews into the special OFT box, the Any-maze software was used for tracking and identification immediately. By tracking the trajectory map, we observed that the tree shrews moved freely in the special OFT box. Each of them ran and jumped freely in line with the basics of the tree shrews' living habits (Figure 2A). Through the comparison of the initial and re-measurement trajectories, the activity patterns of the same tree shrews are similar in the dedicated squatting box, indicating that the tree shrews' special OFT box is relatively stable in detecting tree shrews' activity (Figure 2A).





Figure 2: Partial trajectories of initial and re-measured OFT in tree shrews (A) Representative images of trajectories from 3 tree shrews are illustrated: Test1: trajectories of initial OFT; Test2: trajectories of re-measured OFT; (B and C) Quantitation of the activities of the tree shrews from 2 tests by our OFT system showed the data were stable. Data are presented as the means \pm SEM

The Distance and Average Speed in the Open Field test Experiment have Extremely High Reliability

The intra-group correlation coefficient (ICC) analysis was carried out for the initial distance and the mean speed of the tree shrews in the open field test experiment. The reliability was evaluated by the ICC parameters. The results showed that the activity distance's ICC is 0.951 ($p < 0.001$) and the average speed's ICC is 0.962 ($p < 0.05$) between the first test and the second test. It suggested that the distance and the averaged speed reliability in the open field test experiment were extremely high. Moreover, after paired *t*-test on the above parameters, it was found that the data of the two measurements were stable and there was no significant difference (Figure 2B and C, A: $p = 0.591$; B: $p = 0.685$, $n = 11$).


Stress Significantly Reduced the Activity of the Tree Shrews

To investigate the applicability and generality of our open field test experiment system, the activity of tree shrews in pre-stress (Baseline) and 2 weeks after stress (Stress week 2) in the OFT experiment were observed. As shown in Figure 3, the activity of tree shrews was significantly reduced after light stress for 2 weeks (Figure 3A). Comparing the distance of each tree shrew with its baseline, it was found that the tree shrews' activity distance was significantly decreased after 2 weeks of stress (Figure 3B, $p < 0.01$, $n = 15$). Moreover, we not only observed the alteration of horizontal movement, but also investigated the numbers of jumping in these tree shrews. Among these tree shrews, the numbers of jumping for 12 tree shrews showed a significant decrease (Figure 3C, $p < 0.05$, $n = 12$) while 3 in 15 tree shrews showed the increased jumping (data not shown). All these suggested that 2-week staying up late contributed to the change of the activities in the tree shrews and this has a lot to do with individual differences in tree shrews. Furthermore, to further validate the stress-induced stress behavior tree shrew model, we also collected tree shrew morning urine before and after the stress. We found that the ACTH concentration in the urine of the tree shrews was significantly increased after 2 weeks of stress (Figure 4, $p < 0.001$, $n = 15$). All these data suggesting that our OFT system was stable and sensitive to investigate the alteration of the activities of the tree shrews.



Data are presented as the means \pm SEM. * $p < 0.05$ and ** $p < 0.01$ compared with baseline

Figure 3: After two weeks of stress, the activities of tree shrews were significantly decreased (A) The representative trajectories of tree shrews after 2-week stress are illustrated; (B) The distance of tree shrews was reduced after 2-week light stress; (C) The effects of light stress on the numbers of jumping of tree shrews



After 2-week of light stress, the concentration of ACTH (pg/ml) in the urine of tree shrews in the morning was significantly increased. (***) $p < 0.001$ compared with baseline. Data were normalized to the baseline)

Figure 4: 2-week of light stress enhanced the urine ACTH concentration in tree shrews

Discussion

Physiological and behavioral responses to acute stress can be adaptive, but exposure to chronic stress, especially chronic psychosocial stress, can have negative consequences and increase susceptibility to chronic diseases [15]. Stress refers to the internal reaction of the body to the environment and is caused by the loss and damage of the body caused by the outside world. Classification according to the results of stress may have pathological stress and physiological stress. In recent years, the role of stress-induced neuroendocrine system changes has attracted much attention. The most important pathological change is the excessive drive of the Hypothalamic-Pituitary-Adrenal (HPA) axis [16]. The core driver of the HPA axis activation response is the corticotropin-releasing hormone (CRH) located in the paraventricular nucleus (PVN) site [17]. In human patients, there is a significant increase in neuronal apoptosis in the norepinephrine-blue-spot system and the amine-based system, with the norepinephrine-blue-spot system being the most severely damaged [18]. By placing the animals in a mild stress-stimulated environment, after several weeks of action (typically 2-3 weeks), animals exhibit clinically common symptoms in depressed patients [19]. OFT is a classic method for evaluating the motor function and stress behavior in experimental animals and is widely used in the basic research of neuroscience. In the open field test experiment of tree shrews, the horizontal movement in the ground direction reflects the locomotor activity, and the vertical ground movement including climbing and jumping reflects the curiosity of the tree shrews to the fresh environment. The number of faecal particles reflects the degree of tension of the tree shrews; the number of crossings reflects the anxiety state of the tree shrews, and the stress state can lead to a certain decrease in value. Therefore, the reliability of OFT and the stability of its repeated test parameters are very important [20,21]. In previous experiments, tree shrews' open field test was conducted in a home cage or a rodent open-space box, and they differed greatly from the tree shrews' living habits, with low accuracy and repeatability. Compared with the previous design, we designed the tree shrews' specific OFT box with higher accuracy and repeatability. It is also important to establish a behavioral experimental system with high reproducibility and suitable for tree shrews' habits. Degree refers to the degree of consistency of multiple measurement results, which reflects the stability and reliability of a measurement method or a tool. The test-retest reliability is a kind of reliability, which reflects the measurement method or a tool that consistently measures the consistency of two measurements at different times. The commonly used statistical evaluation parameter for the test-retest reliability is the ICC parameter. The ICC parameter, the intra-group correlation coefficient, was first proposed by Bartko in 1966 for evaluating the reliability of continuous or grade variables [22,23]. If the ICC is greater than 0.75, the credibility is good. If it is between 0.5 and 0.75, the credibility is generally acceptable. And when it is less than 0.5, the credibility is poor. The results of ICC for tree shrews tested in the special OFT box suggested that the common parameters of the active distance and the average speed had extremely high reliability, confirming that the tree shrews' specific OFT system has a good test-retest reliability. The system is stable. Meanwhile, there was no signal loss when we used Any-maze software to track tree shrews in the special OFT box. The trajectory of the tree shrews in the OFT box was well captured. As our data showed, the open field test experiment which is carried out by our special OFT box for tree shrews, restored the basic activities of the tree shrews to a large extent and the data from our OFT system is stable. Moreover, the moving distance and average speed have extremely high reliability in selecting the evaluation parameters. Besides, the results of the tree shrews' urine test showed that ACTH was significantly increased after two weeks of light-stress. Consistent with this, after 2 weeks of stress, the activity of tree shrews by our OFT system changed significantly, that is, the distance of tree shrews was significantly reduced after stress. Taken together, our OFT system can investigate the change of activity of tree shrews after stress.

Conclusion

The tree shrews' specific OFT system that we designed can help us to get the stable and reliable data. Moreover, it is sensitive to test the effect of light-stress on the activities and anxiety of tree shrews, which is consistent with the increased ACTH level after a 2-week staying up late.

Acknowledgment

This work was financially supported by grants from the National Natural Science Foundation of China (81801314), the Natural Science Foundation of the Jiangsu Higher Education Institutions of China (18KJB180016) to W. J, and the Key R&D Program of Jiangsu Province (BE2016761 and 2017CX010) to J.G.. The authors thank all members of each department. The authors thank Mr. Natan for his review of the overall language of the manuscript.

References

1. Fuchs E, Kramer M, Hermes B, Netter P, Hiemke C (1996) Psychosocial stress in tree shrews: clomipramine counteracts behavioral and endocrine changes. *Pharmacol Biochem Behav* 54: 219-28.
2. Kramer M, Hiemke C, Fuchs E (1999) Chronic psychosocial stress and antidepressant treatment in tree shrews: time-dependent behavioral and endocrine effects. *Neurosci Biobehav Rev* 23: 937-47.
3. Lupien SJ, McEwen BS, Gunnar MR, Heim C (2009) Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nat Rev Neurosci* 10: 434-45.
4. Numa C, Nagai H, Taniguchi M, Nagai M, Shinohara R, et al. (2019) Social defeat stress-specific increase in c-Fos expression in the extended amygdala in mice: Involvement of dopamine D1 receptor in the medial prefrontal cortex. *Sci Rep* 9: 16670.
5. Popoli M, Yan Z, McEwen BS, Sanacora G (2011) The stressed synapse: the impact of stress and glucocorticoids on glutamate transmission. *Nat Rev Neurosci* 13: 22-37.

6. Mccarty RJP (2017) Optimizing laboratory animal stress paradigms: The H-H* experimental design. *Psychoneuroendocrinology* 75: 5-14.
7. Fan Y, Yu DD, Yao YG (2014) Tree shrew database (TreeshrewDB): a genomic knowledge base for the Chinese tree shrew. *Scientific Reports* 4: 1426.
8. Magarinos AM, McEwen BS, Flugge G, Fuchs E (1996) Chronic psychosocial stress causes apical dendritic atrophy of hippocampal CA3 pyramidal neurons in subordinate tree shrews. *J Neurosci* 16: 3534-40.
9. Wang J, Xu XL, Ding ZY, Mao RR, Zhou QX, et al. (2013) Basal physiological parameters in domesticated tree shrews (*Tupaia belangeri chinensis*). *Dongwuxue Yanjiu* 34: E69-74.
10. Wang J, Zhou QX, Tian M, Yang YX, Xu L (2011) Tree shrew models: a chronic social defeat model of depression and a one-trial captive conditioning model of learning and memory. *Dongwuxue Yanjiu* 32: 24-30.
11. Wang J, Chai AP, Zhou QX, Longbao LV, Wang L, et al. (2013) Chronic Clomipramine Treatment Reverses Core Symptom of Depression in Subordinate Tree Shrews. *Plos One* 8: e80980.
12. Czeh, B, Fuchs E, Wiborg O, Simon M (2016) Animal models of major depression and their clinical implications. *Prog Neuropsychopharmacol Biol Psychiatry* 64: 293-310.
13. Mineur YS, Belzung C, Crusio WE (2006) Effects of unpredictable chronic mild stress on anxiety and depression-like behavior in mice. *Behav Brain Res* 175: 43-50.
14. Zimprich A, Garrett L, Deussing JM, Wotjak CT, Fuchs H, et al. (2014) A robust and reliable non-invasive test for stress responsivity in mice. *Front Behav Neurosci* 8: 125.
15. Hollis F, Isgor C, Kabbaj M (2013) The consequences of adolescent chronic unpredictable stress exposure on brain and behavior. *Neurosci* 249: 232-41.
16. Porsolt RD (1979) Animal model of depression. *J Biomedicine* 30: 139-40.
17. Abas MJBR, Therapy (1995) *Handbook of affective disorders* (2nd Edn) Churchill Livingstone, Edinburgh, England.
18. Gonzalez MM, Aston-Jones G (2008) Light deprivation damages monoamine neurons and produces a depressive behavioral phenotype in rats. *Proc Natl Acad Sci USA* 105: 4898-903.
19. Willner PJP (1997) Validity, reliability and utility of the chronic mild stress model of depression: a 10-year review and evaluation. *Psychopharmacol* 134: 319-29.
20. Prut L, Belzung C (2003) The open field as a paradigm to measure the effects of drugs on anxiety-like behaviors: a review. *Eur J Pharmacol* 463: 3-33.
21. Ramos A (2008) Animal models of anxiety: do I need multiple tests 29: 493-8.
22. Berk RAJAO (1979) Generalizability of behavioral observations: a clarification of interobserver agreement and interobserver reliability. *Am J Ment Defic* 83: 460-72.
23. Watkins MW, Pacheco MJ (2000) Interobserver Agreement in Behavioral Research: Importance and Calculation. *J Behav Educ* 10: 205-12.

Submit your next manuscript to Annex Publishers and benefit from:

- Easy online submission process
- Rapid peer review process
- Online article availability soon after acceptance for Publication
- Open access: articles available free online
- More accessibility of the articles to the readers/researchers within the field
- Better discount on subsequent article submission

Submit your manuscript at
<http://www.annexpublishers.com/paper-submission.php>